

On Conformal Infinity and Compactifications of the Minkowski Space

Arkadiusz Jadczyk*

*Center CAIROS, Institut de Mathématiques de Toulouse
Université Paul Sabatier, 31062 TOULOUSE CEDEX 9, France*

September 15, 2010

Abstract

Using the standard Cayley transform and elementary tools it is reiterated that the conformal compactification of the Minkowski space involves not only the “cone at infinity” but also the 2-sphere that is at the base of this cone. We represent this 2-sphere by two additionally marked points on the Penrose diagram for the compactified Minkowski space. Lacks and omissions in the existing literature are described, Penrose diagrams are derived for both, simple compactification and its double covering space, which is discussed in some detail using both the $U(2)$ approach and the exterior and Clifford algebra methods. Using the Hodge \star operator twistors are realized as spinors (i.e., vectors of a faithful irreducible representation of the even Clifford algebra) for the conformal group $O(4, 2)$. Killing vector fields corresponding to the left action of $U(2)$ on itself are explicitly calculated. Isotropic cones and corresponding projective quadrics in pseudo-Hermitian spaces $H_{p,q}$ are also discussed. Applications to flat conformal structures, including the normal Cartan connection and conformal development has been discussed in some detail.

*E-mail address: arkadiusz.jadczyk@cict.fr

1 Introduction

The term *compactification* can have several different meanings. Given a manifold \mathcal{M} we may try to embed it into a compact one and take its closure. Or, we can attach to \mathcal{M} ideal boundary points or boundary components so as to obtain a compact space. In physics compactification of space–time can be used either in order to study its conformal invariance, or to study its asymptotic flatness, or its singularities. In the available literature the differences between these different approaches are not always made clear and the mathematical language involved is not always as precise as one would wish.

This paper is a compromise between being completely self–contained and a typical specialized article. We use techniques of algebra and geometry but we avoid twistor notation of Penrose school which can be confusing to many mathematicians. The paper is aimed at mathematicians interested in mathematical properties of Minkowski space related to projective geometry, and at mathematical physicists interested in the subject. Relativists will find next to nothing of interest for them in the material below (perhaps except of a warning about how errors can easily propagate). They have their own aims and techniques and, as a rule, are usually not interested in generalizations going beyond four space–time dimensions.

In section 2 we review the conformal compactification $\tilde{M} = U(2)$ of the Minkowski space M . We are following there the elegant and simple method of A. Uhlmann [1] by using 2×2 matrices and the Cayley transform. We are also investigating in some detail the structure of the “light cone at infinity”, that is the set difference $\tilde{M} \setminus M$ and point out that it consists not only of the (double) light cone, but also of a 2-sphere that connects the two cones - a fact that was known to Roger Penrose [2, p. 178]. This fact was not always realized by other authors writing on this subject even when they quoted Penrose (cf. e.g., Sec. 3). Additionally, as a complement to this particular representation of \tilde{M} , in appendix A, we calculate vector fields on M corresponding to one–parameter subgroups of $U(2)$ acting on itself by left translations.

In section 3, as an educational example, we discuss in some detail the faulty argument and the missing 2-sphere in [3]. In particular we reproduce a crucial part of reasoning used in [3] and point out the omission explicitly. Similar omissions, this time taken from [14] and also from a recent papers on conformal field theory, are discussed in section 3.2.

In section 4, geometrical representation of the conformal compactification \tilde{M} is discussed using the cylinder representation of Einstein's static universe - the standard representation in general relativity. This leads to a two-dimensional diagram - a version of the Penrose diagram (cf. Fig. 1), with the two 2-spheres that need to be identified. Owing to this identification no intrinsic distinction between \mathcal{J}^+ and \mathcal{J}^- is possible. In Fig. 3 and Fig. 6 we mark these two parts of the conformal infinity in order to be able to compare this diagram with those (as in Fig. 4) found in the standard literature.

In section 5, the explicit action of the Poincaré group on the conformal infinity is calculated, where it is in particular shown that this action is transitive there. A lack of a mathematical precision in the mathematical literature on the subject is also elucidated.

Section 6 starts with a simple exercise showing a geometrically amusing fact that null geodesics can be completely trapped at infinity. A role of the conformal inversion, and the signature of the induced metric is also discussed there. Then, a pictorial representation of the infinity is given, first as a double cone with identified vertices in Fig. 3, then, more correct as far as its differentiability properties are concerned, as a squeezed torus in Fig. 6. A typical, almost identically looking, but with a different meaning, picture - taken from [4] - is shown in Fig. 4. The squeeze point in Fig. 6 corresponds to what is usually denoted as I^0, I^+, I^- (or i^0, i^+, i^-) in the standard literature. All three points coincide in our case.¹ A correct image, which we reproduce here in Fig. 5 can be found in Fig. 2 of [5]. It may be worth quoting the following remarks from the monograph of Penrose and Rindler [2, p. 298]:

“Having this natural association between the points of \mathcal{J}^- and \mathcal{J}^+ , for Minkowski space, it is in some respect natural to make identification between \mathcal{J}^- and \mathcal{J}^+ , the point A^- being identified with A^+ and \mathcal{J}^- and \mathcal{J}^+ written as \mathcal{J} . If we do this, then, for the sake of continuity we should also identify I^- with I^0 , and I^0 with I^+ .”

To which they added:

“For reasons that we shall see in more detail later, such identification cannot be satisfactorily carried out in *curved* asymptotically flat spaces. (Not only is there apparently no *canonical*

¹A. Uhlmann [1] conjectured that it may be a squeezed Klein's bottle. Klein's bottle is unnecessary as long as we do not care about the embedding. Squeezed torus does the job.

way of performing such identifications in general, but, when the total mass is non-zero *any* identification would lead to failure of the required regularity conditions along the identification hypersurface.) For many purposes, the identification of \mathcal{J}^- with \mathcal{J}^+ may, even in Minkowski space, seem unphysical (and, of course, it need not be made). However, for various mathematical purposes the identification is very useful...”

In subsection 6.4 we discuss the double cover of \tilde{M} , that can be obtained by the same method as in section 3 but by considering positive rays rather than generator lines.² This leads us to the compactification with the past infinity \mathcal{J}^- and future infinity \mathcal{J}^+ different, but I^- and I^+ are identified, though different from I^0 . The resulting Penrose diagram is given in Fig. 2, and the ensuing graphic representation of the conformal infinity is pictured in Fig. 7 and in Fig. 8.

We follow here method used by Kopczyński and Woronowicz in [9], but this time applied to the double cover of M . Moreover, we identify the antilinear map $x \mapsto x^\perp$ used by these authors as a Hodge \star operator adapted for a complex vector space V equipped with a non-degenerate sesquilinear form³. After a general introduction, for an arbitrary signature, starting with the Grassmann algebra endowed with the natural scalar product, we specialize to the case of signature $(2, 2)$, $V \approx H_{2,2}$, and relate the two compactification methods - one in which the points of the double covering of the compactified Minkowski space are represented by oriented maximal isotropic subspaces of a four dimensional complex space endowed with a sesquilinear form of signature $(2, 2)$, and the one discussed in Sec. 6.4 based on rays of the null cone in 6-dimensional real space endowed with a scalar product with signature $(4, 2)$. We derive explicit formulas connecting the $U(2)$ compactification and the one based on $H_{2,2}$.

In order to show how the compactified Minkowski space enters more general conformal structures on manifolds, in section 8 we briefly review geometry of conformal structures, second-order frames and the normal Cartan connection. We end this section by explicitly calculating the standard embedding of Minkowski space into the compact projective hyperquadric using

²This construction is also briefly mentioned in [6, p. 180]. It is also worthwhile to mention that $(U(1) \times SU(2))/Z_2$, $Z_2 = \{I, -I\}$, with the topology of $(S^1 \times S^3)/Z_2$ is homeomorphic, as a manifold, to its double cover $U(1) \times SU(2)$ - cf. [7] and [8].

³For a discussion in case of positive definite scalar product cf. e.g., [10].

the conformal development.

Appendix B can be read independently of the rest of the paper. While in section 6 we have made use of the fact that, as discussed in some detail in [9], there is a one-to-one correspondence between null geodesics in \tilde{M} and isotropic lines in $V \approx H_{2,2}$, in appendix B the structure of this set is studied for a general case of signature (p, q) .

2 Conformally compactified Minkowski space

In this section we follow idea of Armin Uhlmann [1]. Let $H(2)$ be the real vector space of complex 2×2 Hermitian matrices. Let M be the Minkowski space endowed with the standard coordinates x^1, x^2, x^3, x^0 ,⁴ and the quadratic form $q(x) = -(x^0)^2 + (x^1)^2 + (x^2)^2 + (x^3)^2$, and let $\varphi : M \rightarrow H(2)$ be the isomorphism given by⁵

$$\varphi(x) = X = \begin{bmatrix} x^0 + x^3 & x^1 - ix^2 \\ x^1 + ix^2 & x^0 - x^3 \end{bmatrix}. \quad (1)$$

Then we have

$$\det(X) = (x^0)^2 - ((x^1)^2 + (x^2)^2 + (x^3)^2) = -q(x). \quad (2)$$

Let $U(2)$ be the group of all unitary 2×2 matrices with complex entries. Let $u : H(2) \rightarrow U(2)$ be the Cayley transform:

$$u(X) = U = \frac{X - iI}{X + iI}.$$

Notice that, because of X being Hermitian, $\det(X + iI) \neq 0$. We then have

$$\begin{aligned} I + U &= \frac{iI + X + X - iI}{X + iI} = \frac{2X}{X + iI}, \\ I - U &= \frac{X + iI - X + iI}{X + iI} = \frac{2i}{X + iI}. \end{aligned} \quad (3)$$

In particular $\det(I - U) \neq 0$ and

$$X = i \frac{I + U}{I - U}. \quad (4)$$

⁴Sometimes, as an alternative, will set $x^0 = x^4$, and write $x = (x^1, \dots, x^4) \in M$.

⁵Cf. e.g., [11, p. 324].

It easily follows that $\psi = u \circ \varphi : M \rightarrow U(2)$ is a bijection from M onto the open subset of $U(2)$ consisting of those U for which $\det(I - U) \neq 0$.

Remark 1. *It may be useful for the reader to see the explicit form of $\psi(x)$ for any $x \in M$, namely*

$$U = \psi(x) = \frac{1}{-q(x) - 1 + 2ix^0} \begin{bmatrix} 1 - q(x) + 2ix^3 & 2(ix^1 + x^2) \\ 2(ix^1 - x^2) & 1 - q(x) - 2ix^3 \end{bmatrix}. \quad (5)$$

We also have, explicitly:

$$\det(I - U) = \frac{4}{1 + q(x) - 2ix^0}, \quad \det(I + U) = \frac{4q(x)}{1 + q(x) - 2ix^0}. \quad (6)$$

The first one of the last two equalities shows that for any $U \in \psi(M)$, $\det(I - U) \neq 0$, while the second one states that $\det(I + U) = 0$ if and only if $q(x) = 0$. Notice that the quantity $1 + q(x) - 2ix^0 \neq 0$ for all $x \in M$.

Let us now determine the structure of the remaining set \mathfrak{J} :

$$\mathfrak{J} = U(2) \setminus \psi(M) = \{U \in U(2) : \det(I - U) = 0\}.$$

Let $m : U(2) \rightarrow U(2)$ be the diffeomorphism of $U(2)$ given by

$$m(U) = -U,$$

i.e., the group translation by $-I$. Let us investigate the structure of the set $m(\mathfrak{J})$ - the image of $\mathfrak{J} \subset U(2)$ under m . We split this set into two disjoint non empty components \mathfrak{J}_c and \mathfrak{J}_s defined by

$$\mathfrak{J}_c = m(\mathfrak{J}) \setminus \mathfrak{J}, \quad \text{and} \quad \mathfrak{J}_s = m(\mathfrak{J}) \cap \mathfrak{J}.$$

Remark 2. *To see that both sets, \mathfrak{J}_c and \mathfrak{J}_s , are non empty, notice that $U_0 = -I = m(I)$ is not in \mathfrak{J} , but is in $m(\mathfrak{J})$. Therefore U_0 is in \mathfrak{J}_c . On the other hand let $U_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Then U_1 and $-U_1 = m(U_1)$ are in \mathfrak{J} , thus U_1 is in \mathfrak{J}_s .*

The set \mathfrak{J}_c is, by its definition in the range of Cayley transform, therefore we can apply ψ^{-1} to \mathfrak{J}_c .

Denoting by K the light cone through the origin:

$$K = \{x \in M : q(x) = 0\},$$

let us show that

$$\psi^{-1}(\mathfrak{J}_c) = K. \quad (7)$$

With $x \in M$ we have that $x \in \psi^{-1}(\mathfrak{J}_c)$ if and only if $\psi(x) \in \mathfrak{J}_c$, that is if and only if $(U \in m(\mathfrak{J}))$ and $(U \notin \mathfrak{J})$. That is $x \in \psi^{-1}(\mathfrak{J}_c)$ if and only if $\det(I + U) = 0$ and $\det(I - U) \neq 0$. It follows now from Eq. (6) that $\det(I - U)$ is automatically non-zero, and that $\det(I + U) = 0$ is equivalent to $q(x) = 0$, that is $x \in K$.

It remains to identify the set \mathfrak{J}_s . Let $j : U(2) \rightarrow U(2)$ be the map

$$j(U) = iU,$$

i.e., the translation by i . It follows from the very definition that $U \in \mathfrak{J}_s$ is equivalent to: $\det(I - U) = 0$ and $\det(I + U) = 0$. It follows that $U \in \mathfrak{J}_s$ if and only if one eigenvalue of U is equal $+1$ while the other eigenvalue is equal -1 . It follows that $j(U) = iU$ has eigenvalues $+i$ and $-i$. Therefore $I - iU$ is invertible and $U = \varphi(X)$, with X given by Eq. (4) and U replaced by iU . It follows that $j(U)$ is in the range of ψ . Thus we conclude that $j(\mathfrak{J}_s) \subset \psi(M)$. Let us show that $\psi^{-1}(j(\mathfrak{J}_s))$ is the 2-sphere:

$$\psi^{-1}(j(\mathfrak{J}_s)) = \{x \in M : x^0 = 0, (x^1)^2 + (x^2)^2 + (x^3)^2 = 1\}.$$

With $U \in \mathfrak{J}_s$ let $x = \psi^{-1}(j(U))$. Then

$$\psi(x) = X = i \frac{I + iU}{I - iU}.$$

It follows that X has eigenvalues $i \frac{1+i}{1-i} = -1$ and $i \frac{1-i}{1+i} = 1$, which is equivalent to $\det(X) = -1$ and $\text{tr}(X) = 0$. Now, from Eq. (1) it follows that $\text{tr}(X) = 0$ is equivalent to $x^0 = 0$, and then $\det(X) = -1$ is equivalent to $(x^1)^2 + (x^2)^2 + (x^3)^2 = 1$, which concludes our proof.

It follows from the above that $U(2) \setminus \psi(M)$ consists of two pieces. The first piece is the set of all unitary matrices with precisely one eigenvalue equal to -1 , the other eigenvalue different from $+1$. This piece has the structure of the *light cone at infinity*. The matrix $U = -I$ is the apex of this cone. The second piece consists of unitary matrices with one eigenvalue equal to -1 , the other eigenvalue being $+1$. This piece is the *2-sphere at infinity* that forms “a base” of the light cone at infinity.

Remark 3. *A closely related derivation of this fact can be found in [12, Theorem 6]. This pedagogical paper is closely related in spirit and is a recommended reading for all those interested in the subject.*

Remark 4. *It is easy to calculate the result of the transformation $x \mapsto x'$ corresponding to the left translation $U \mapsto iU = j(U)$. The result of a simple calculation reads:*

$$x^{0'} = \frac{1 + q(x)}{1 - q(x) - 2x^0} \quad \text{and} \quad \mathbf{x}' = \frac{2\mathbf{x}}{1 - q(x) - 2x^0}.$$

This particular transformation can be interpreted in terms of conformal transformations $T(a)x = x + a$, $K(a) = RT(a)R$, $D(\lambda)x = \lambda x$, where R is the inversion $R(x) = x/q(x)$. A simple calculation shows that

$$x' = T(-a)D(2)K(a)x,$$

where $a^0 = -1$, $\mathbf{a} = 0$. The transformation is singular on the light cone centered at $-a$.

In appendix A we calculate the conformal vector fields on Minkowski space corresponding to left actions of one-parameter subgroups of $U(2)$.

3 The overlooked 2-sphere

In their *Introduction to Twistor Theory* [3, Chapt. 5], *Compactified Minkowski Space*, the authors obtain their “cone at infinity” using a different method and, as we will see, their incomplete reasoning leads to their neglecting of the 2-sphere at infinity. First, we will reproduce their reasoning, using their notation, with slight changes, simplifications, and with some elucidating comments. Then, we will present our corrected derivation and its result.

3.1 Reasoning of Huggett and Tod

Here we will present the essence of the reasoning in [3], though with some changes of the notation. We denote by M the standard Minkowski space, that is $E^{3,1} = \mathbb{R}^3 \oplus \mathbb{R}^1$, with coordinates $x = (\mathbf{x}, t)$, endowed with the quadratic form $q(x) = \mathbf{x}^2 - t^2$, where $\mathbf{x} = (x^1, x^2, x^3)$, and \mathbf{x}^2 is the standard Euclidean quadratic form of \mathbb{R}^3 : $\mathbf{x}^2 = (x^1)^2 + (x^2)^2 + (x^3)^2$. Let $E^{1,1}$ be \mathbb{R}^2 endowed with the quadratic form q_2 defined by $q_2(x_5, x_6) = (x^5)^2 - (x^6)^2$, $(x_5, x_6) \in \mathbb{R}^2$. We denote by $E^{4,2}$ the 6-dimensional space $E^{3,1} \oplus E^{1,1}$, with coordinates $(Z^\alpha) = (x, x_5, x_6)$, and endowed with the quadratic form $Q(x, x_5, x_6) = q(x) + q_2(x_5, x_6)$. In order to simplify a bit the notation, let us set, in this section,

$$x^5 \equiv v, \quad x^6 \equiv w.$$

Let \mathcal{N} be the null cone of $E^{4,2}$ minus the origin:

$$\mathcal{N} = \{Z \in E^{4,2} : Z \neq 0 \text{ and } Q(Z) = 0\}, \quad (8)$$

and let $P\mathcal{N}$ be the set of its generators, that is the set of straight lines through the origin in the directions nullifying $Q(Z)$. In other words $P\mathcal{N} = \mathcal{N} / \sim$, where, for $Z, Z' \in \mathcal{N}$, $Z \sim Z'$ if and only if there exists a nonzero $\mu \in \mathbb{R}$ such that $Z' = \mu Z$. We denote by π the projection $\pi : \mathcal{N} \rightarrow P\mathcal{N}$. Then $P\mathcal{N}$, with its projective topology, is a compact projective quadric. $P\mathcal{N}$ is called the *compactified Minkowski space*.

Consider now the following smooth map between manifolds: $\tau : M \rightarrow E^{4,2}$ given by the formula:

$$\tau(\mathbf{x}, t) = (\mathbf{x}, t, \frac{1}{2}(1 - q(x)), -\frac{1}{2}(1 + q(x))). \quad (9)$$

The map τ is evidently injective. Let \mathcal{Z} be the hyperplane in $E^{4,2}$:

$$\mathcal{Z} = \{Z \in E^{4,2} : v - w = 1\}. \quad (10)$$

Lemma 1. *The image $\tau(M)$ in $E^{4,2}$ coincides with the intersection $\mathcal{N} \cap \mathcal{Z}$ of \mathcal{N} with \mathcal{Z} .*

Proof. It is clear that $\tau(x) \neq 0$, and it also follows by an easy calculation that $Q(\tau(x)) = 0$. Evidently, from Eq. (9), $\tau(x)$ is also in \mathcal{Z} . Conversely, let $Z = (x, v, w)$ be in $\mathcal{N} \cap \mathcal{Z}$. From $Q(Z) = Q(x, v, w) = 0$ we get $q(x) + v^2 - w^2 = 0$. But $v^2 - w^2 = (v - w)(v + w)$ so that from $v - w = 1$ it follows that $q(x) + v + w = 0$. Together with $v - w = 1$ it implies $q(x) + 2v = 1$ or $v = \frac{1}{2}(1 - q(x))$ and $w = -\frac{1}{2}(1 + q(x))$. It follows that $Z = \tau(X)$. \square

From now on we will follow the arguments in [3, p. 36] step by step, skipping what is not essential and adapting to our notation.

“On any generator of \mathcal{N} with $v - w \neq 0$, we can find a point satisfying $v - w = 1$ and hence a point in M . Thus M is identified with a subset of $P\mathcal{N}$.”

This is clear. If (x, v, w) is in \mathcal{N} and $v - w \neq 0$, then $\frac{x}{v-w}$ is in $\mathcal{N} \cap \mathcal{Z}$.

“The points in $P\mathcal{N}$ not in $\tau(M)$ corresponds to the generators of $P\mathcal{N}$ with $v - w = 0$.”

This is evident from the definition. Now there comes an unclear paragraph with an erroneous conclusion:

“This is the intersection of N with a null hyperplane through the origin. All such hyperplanes are equivalent under $O(4, 2)$ so to see what these extra points represent, we consider the null hyperplane $v + w = 0$. From Eq. (9) we see that the points of M corresponding to generators of \mathcal{N} which lie in this hyperplane are just the null cone of the origin. Thus $P\mathcal{N}$ consists of $\tau(M)$ with an extra cone at infinity.”

It is rather hard to follow this fuzzy reasoning, therefore we will study the structure of the “extra part” directly from the definition. The extra part is the projection by π of those points in \mathcal{N} for which $v - w = 0$. Now the following two cases must be considered separately: either $v = w = 0$ or $v = w \neq 0$. Let $\mathcal{N}_c = \{Z \in \mathcal{N} : v = w \neq 0\}$, and $\mathcal{N}_s = \{Z \in \mathcal{N} : v = w = 0\}$. Each element of $\pi(\mathcal{N}_c)$ has a unique representative $Z' = (x', v', w')$ in \mathcal{N} with $v' = w' = 1$. Since $Q(Z') = 0$, we have $q(x') = 0$. Therefore $\pi(\mathcal{N}_c)$ has the structure of the null cone at zero in M . But there is also the second part, $\pi(\mathcal{N}_s)$. If $Z = (\mathbf{x}, t, 0, 0)$ is in \mathcal{N}_s , then $t \neq 0$, otherwise, because of $Q(Z) = q(x) = 0$ we would have $\mathbf{x} = 0$. Therefore each $Z = (\mathbf{x}, t, 0, 0)$ in \mathcal{N}_s has a unique representative with $t = 1$. From $q(x) = 0$ it follows then that $\mathbf{x}^2 = 1$. It follows that $\pi(\mathcal{N}_s)$ has the structure of the 2-sphere. This part is missing in the conclusion of [3]. One of the possible reasons for this omission can be a possibly misleading statement in Penrose and Rindler [2, p. 303], where we can read

“... and the remainder of the intersection of the 4-plane with \tilde{M} is \mathcal{J} (the identified surfaces $\mathcal{J}^+, \mathcal{J}^-$ of the previous construction).”

The point is that in \mathcal{J} of Penrose and Rindler one has to first identify the two 2-spheres, one of \mathcal{J}^+ and one of \mathcal{J}^- , though with opposite orientations - see the next subsection. This lack of precision in [2] may have confused the authors of [3, 13].

3.2 The 2-sphere missed by Akivis and Goldberg

A similar inadvertency takes place in a monograph on conformal geometry by M. A. Akivis and V. V. Goldberg [14]. In the introductory chapter the

authors analyze the Euclidean case. They start with the equation of a hypersphere in the conformal space C^n , which is just $E^{n,0}$ endowed with an Euclidean scalar product defined up to a non-zero multiplicative constant. The equation, in *polyspherical coordinates* s^0, s^i, s^{n+1} , reads:

$$s^0 \sum_{i=1}^n (x^i)^2 + 2 \sum_{i=1}^n s^i x^i + 2s^{n+1} = 0.$$

When $s^0 \neq 0$, this can be put in the form:

$$\sum_{i=1}^n (x^i - a^i)^2 = r^2,$$

where

$$a^i = -\frac{s^i}{s^0}, \quad r^2 = \frac{1}{(s^0)^2} \left(\sum_{i=1}^n (s^i)^2 - 2s^0 s^{n+1} \right).$$

In order to describe a hypersphere of zero radius (centered at a^i) we must have

$$(X, X) := \sum_{i=1}^n (s^i)^2 - 2s^0 s^{n+1} = 0,$$

which is just the equation (8) of the null cone \mathcal{N} in $E^{n+1,1}$ with $s^0 = \frac{1}{2}(w - v)$, $s^{n+1} = (w + v)$, adapted to the Euclidean signature. Hyperspheres of zero radius correspond to the points (their centers) in C^n . The remaining set of non-zero solutions of Eq. (8) is the line $s^0 = 0$, $x^i = 0$, $1, \leq i \leq n$, $s^{n+1} \neq 0$ - the point at infinity.

The same strategy is then used in Chapter 4.1 in the pseudo-Euclidean case. With slight changes of the notation the authors state [14, p. 127] that

“... after compactification the tangent space $T_x(M)$ is enlarged by the point at infinity y with coordinates $(0, 0, \dots, 0, 1)$ and by the isotropic cone C_x , with vertex at this point y whose equation is the same as the equation of the cone C_x , namely $g_{ij}x^i x^j = 0$.”

There is a subtle inadvertency there. The change of notation is not important, so let us use the same notation as in the Euclidean case. When $s^0 \neq 0$ we have the same situation as in the Euclidean case, except that the “hypersphere of zero radius” becomes now a cone (light cone in the Minkowski

case). It remains to consider the case of $s^0 = 0$. Here we have two possibilities: either $s^{n+1} = 0$ or $s^{n+1} \neq 0$. If $s^{n+1} = 0$, then, necessarily, the n -vector $(s^i) \neq 0$, and $g_{ij}s^i s^j = 0$. But then, we should consider the set of lines and not the set of points.

For example in the case of Minkowski space we find that the set of lines is the quadric (S^2) , and **not** the “isotropic cone”, as falsely stated in [14]. On the other hand, if $s^{n+1} \neq 0$, then we can choose $s^{n+1} = 1$. In this case no freedom of choosing the scale remains and we get $g_{ij}s^i s^j = 0$ - the isotropic cone, including its origin.

Another mistake takes place during the discussion of the conformal inversion in [14, p. 15-16]. The authors state that

“In the pseudo-Euclidean space \mathbb{R}_q^n , the inversion in a hypersphere S with center at a point A is defined exactly in the same manner as it was defined in the Euclidean space \mathbb{R}^n (...). However, in contrast to the space \mathbb{R}^n , under an inversion in the space \mathbb{R}_q^n not only does the center a of the hypersphere S not have an image but also points of the isotropic cone C_x with vertex at the point a does not have images. To include these points in the domain of the mapping defined by the inversion in \mathbb{R}_q^n , we enlarge the space \mathbb{R}_q^n not only by the point at infinity, ∞ , corresponding to the point a but also by the isotropic cone C_∞ with the vertex at this point. The manifold obtained as the result of this enlargement is denoted by C_q^n :

$$C_q^n = \mathbb{R}_q^n \cup \{C_\infty\}$$

and is called a *pseudoconformal sphere of index q* . (...) Just like conformal space C^n , the pseudoconformal space C_q^n is homogeneous.”

Adding the image of the isotropic cone under inversion does not result in the homogeneous space. In section 6.1 we show that the conformal inversion with respect to the isotropic cone C_0 centered at the origin $0 \in M$ is implemented by the map $(x, v, w) \mapsto (x, -v, w)$. Using the embedding $\tau : M \rightarrow E^{4,1}$ given by Eq. 9 we find that the image of C_0 consists of vectors of the form $(x, \frac{1}{2}, -\frac{1}{2})$ and therefore $\tau(C_0)$ consists of vectors of the form $(x, -\frac{1}{2}, -\frac{1}{2})$. Let now x be a nonzero vector in C_0 , and let a be a vector satisfying $x \cdot a = \frac{1}{2}$. The action of the translation group is given by Eq. (15). It is clear that after translation

by a the point $(x, -\frac{1}{2}, -\frac{1}{2})$ is mapped to $(x, 0, 0)$, which is **not** in the image of C_0 under inversion. Therefore the statement in [14] that adding just the image of C_0 under inversion gives a homogeneous space is erroneous. It is necessary to add the missing sphere.

A similar misleading statement can be found in a paper by N. M. Nikolov and I. T. Todorov in [15], where the authors state that “The points at infinity in \bar{M} form a $D - 1$ dimensional cone with tip at p_∞ , quoting Penrose [18], and then state that “... the Weyl inversion ... interchanges the light cone at the origin with the light cone at infinity”.⁶

4 From Einstein’s static universe to PN

The group $U(1)$ can be identified with the group of complex numbers $z \in \mathbb{C}$ with $|z| = 1$, and the group $SU(2)$ can be thought of as the group of unit quaternions $\{q = v + x\mathbf{i} + y\mathbf{j} + z\mathbf{k} \in \mathbb{H} : |q|^2 = v^2 + \mathbf{x}^2 = 1\}$. Let $E^{4,1}$ denote \mathbb{R}^5 , with coordinates (\mathbf{x}, v, ψ) , and endowed with the quadratic form $q_5(\mathbf{x}, v, \psi) = \mathbf{x}^2 + v^2 - \psi^2$. Writing $z = e^{i\psi}$, we can then represent the group $U(1) \times SU(2)$ (topologically $S^1 \times S^3$) as the cylinder \mathcal{K} in $E^{4,1}$:

$$\mathcal{K} = \{(\mathbf{x}, v, \psi) : \mathbf{x}^2 + v^2 = 1, \psi \in [-\pi, \pi)\}.$$

Lemma 2. *With $E^{4,2}$ endowed with the coordinates $Z = (\mathbf{X}, T, V, W)$, as in the previous section (but we will use capital letters here) let $\lambda : E^{4,1} \rightarrow E^{4,2}$, be the map*

$$\lambda : (\mathbf{x}, v, \psi) \mapsto (\mathbf{X}, T, V, W) = (\mathbf{x}, \sin(\psi), v, -\cos(\psi)). \quad (11)$$

Then $\pi \circ \lambda$ restricted to \mathcal{K} , is $2 : 1$ and surjective: $\pi \circ \lambda(\mathcal{K}) = PN$. Given any two points $(\mathbf{x}_1, v_1, \psi_1)$ and $(\mathbf{x}_2, v_2, \psi_2)$ in \mathcal{K} , we have $\pi \circ \lambda(\mathbf{x}_1, v_1, \psi_1) = \pi \circ \lambda(\mathbf{x}_2, v_2, \psi_2)$ if and only if the following conditions (i-iii) hold

$$(i) |\psi_2 - \psi_1| = \pi, \quad (ii) \mathbf{x}_2 = -\mathbf{x}_1, \text{ and } \quad (iii) v_2 = -v_1.$$

⁶In a private exchange one of the authors (N.M.N) explained to me that the precise statement should read: “The Weyl inversion ... interchanges the **compact** light cone at the origin with the compact light cone at infinity, where the compact light “cone” with a tip at p is defined as $\{q \in \tilde{M} : p \text{ and } q \text{ are mutually isotropic}\}$.” These concepts have been described in [16, Appendix A,C] and [17], and will be developed in their future paper.

Proof. The proof is evident after noticing that $Q(Z) = \mathbf{X}^2 - T^2 + V^2 - W^2 = 0$ can be written as $\mathbf{X}^2 + V^2 = T^2 + W^2$, and, if $Z \neq 0$, then $V^2 + \mathbf{X}^2 > 0$. Therefore on each generator line of \mathcal{N} there are exactly two points $Z, -Z$, with $V^2 + \mathbf{X}^2 = 1$. \square

In order to be able to represent $P\mathcal{N}$ graphically, on a plane, let us introduce the map $\rho : \mathcal{K} \rightarrow [0, \pi] \times [-\pi, \pi] \subset \mathbb{R}^2$ given by

$$\rho : (\mathbf{x}, v, \psi) \mapsto (\xi = \arccos(v), \psi).$$

In Figure 1 the resulting “Penrose diagram” is shown, using the notation




Figure 1: The “Penrose diagram” of Minkowski space.

as in [19, p. 919], but with two distinguished points denoted as S . In this realization they represent one and the same 2-sphere - they need to be identified. The region inside the triangle with vertices at $(0, -\pi), (0, +\pi), (\pi, 0)$ corresponds to the points in the Minkowski space. In order to understand this correspondence, let us first notice that owing to the equation $v^2 + \mathbf{x}^2 = 1$, we have the following relations:

$$\mathbf{X} = \mathbf{x}, T = \sin(\psi), V = \cos(\xi), W = -\cos(\psi), |\mathbf{X}| = \sin(\xi).$$

When $V - W \neq 0$, we get the corresponding point in Minkowski space with coordinates (\mathbf{r}, t) given by the formulae:

$$\begin{aligned} \mathbf{r} &= \frac{\mathbf{X}}{V - W} = \frac{\mathbf{x}}{\cos(\psi) + \cos(\xi)} \\ t &= \frac{T}{V - W} = \frac{\sin(\psi)}{\cos(\psi) + \cos(\xi)} \\ r &= |\mathbf{r}| = \frac{\sin(\xi)}{\cos(\psi) + \cos(\xi)}. \end{aligned}$$

Now, by elementary trigonometric identities we have that:

$$\begin{aligned} \tan\left(\frac{\psi + \xi}{2}\right) &= \frac{\sin(\psi) + \sin(\xi)}{\cos(\psi) + \cos(\xi)}, \\ \tan\left(\frac{\psi - \xi}{2}\right) &= \frac{\sin(\psi) - \sin(\xi)}{\cos(\psi) + \cos(\xi)}. \end{aligned}$$

It follows that

$$t + r = \tan\left(\frac{\psi + \xi}{2}\right), \quad t - r = \tan\left(\frac{\psi - \xi}{2}\right),$$

which are exactly the equations in [19, p. 919], and in [20, p. 121] (with our ψ, ξ corresponding to their t', r' resp.). Each point in the interior of the triangle represents a 2-sphere at time t and radius $r > 0$ centered at the origin of \mathbf{x} -axes. Each points on the open interval $\xi = 0, |\psi| < \pi$ represents the origin ($t = 0, \mathbf{r} = \mathbf{0}$) of the Minkowski coordinate system. The points I^- and I^+ , with $\xi = 0, \psi = \pm\pi$ both correspond to $V = 1, W = 1, T = 0, \mathbf{X} = 0$ - a single point in the compactified Minkowski space, the apex of the null cone \mathcal{N}_c at infinity. Each point of the open intervals \mathcal{J}^\pm corresponds to a 2-sphere $V = W \neq 0, T \neq 0, \mathbf{X}^2 = T^2$. These 2-spheres build \mathcal{N}_c except of its apex $I^+ = I^-$. The point I^0 represent the same point of the compactified Minkowski space as I^\pm . What is misleading in all the standard literature describing the conformal infinity is the neglecting the fact that there are two exceptional points of the diagram, denoted here as I^1 , and corresponding to the parameter values $\xi = \pi/2, \psi = \pm\pi/2$. These two points correspond to $V = W = 0, T = \pm 1, \mathbf{X}^2 = 1$ which is the sphere \mathcal{N}_s discussed at the end of the previous section. These two exceptional points should be identified in order to give the complete representation of the conformal infinity - compare the discussion of these issues in the papers of Roger Penrose [18, 21].

5 Action of the inhomogeneous Lorentz group $ISO(3, 1)$

5.1 Action of $SO(3, 1)$

The homogeneous Lorentz group $SO(3, 1)$ maps the conformal infinity into itself. It is thus of interest to analyze this action in some details. We will show that there are two invariant submanifolds for this action, one consisting of a point, and one being the 2-sphere \mathcal{N}_s . To this end will use the results of W. Rühl [22]. According to [22], his Eqs. (2.18), (2.19), the homogeneous Lorentz group is represented by $SU(2, 2)$ matrices $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ of the form $A = D = \frac{1}{2}(R + (R^*)^{-1})$ and $B = C = \frac{1}{2}(-R + (R^*)^{-1})$ where $\det(R) = 1$ and $*$ denotes Hermitian conjugation. We need to consider two cases: when R is unitary (pure rotations), and when R is Hermitian (pure boosts). In the case of pure rotations we have $R = R^{*-1}$, Therefore, in this case, $A = D = R, B = C = 0$, and the fractional linear action of $SU(2, 2)$ on $U(2)$ becomes:

$$U' = RUR^{-1}.$$

It is clear that the point at infinity corresponding to $U = E$ is invariant. Also the spectrum of U is an invariant of this transformation, therefore the 2-sphere \mathcal{N}_s corresponding to U with eigenvalues ± 1 is mapped into itself. Now consider the boosts, with $R = R^*$. Denote $R_+ = R + (R^*)^{-1}, R_- = -R + (R^*)^{-1}$. The fractional linear transformation corresponding to the boosts are then of the form:

$$U' = (R_+U + R_-)(R_-U + R_+)^{-1}. \quad (12)$$

Evidently the point $U = E$ is left invariant. Consider now the 2-sphere \mathcal{N}_s corresponding to the unitary operators U with eigenvalues ± 1 . These points are characterized by the property $U^2 = E$. Therefore we can rewrite the Eq. (12) as

$$(R_+U + R_-)((R_- + R_+U)U)^{-1} = ZUZ^{-1},$$

where $Z = R_+U + R_-$. It follows that then also $(U')^2 = E$, therefore the Lorentz boosts map the 2-sphere \mathcal{N}_s onto itself. Thus \mathcal{N}_s is an $O(3, 1)$ -invariant submanifold of the conformal infinity.

5.2 Action of the translations

Consider the translation by a four-vector $a \in M$. Using Clifford algebra methods and the formula for the translations in [23, p. 87] it is easy to calculate the effect of the translation in terms of variables (x, v, w) of section 3.1:

$$x' = x - (v - w)a \quad (13)$$

$$v' = v + (x \cdot a) - \frac{a^2}{2}(v - w) \quad (14)$$

$$w' = w + (x \cdot a) - \frac{a^2}{2}(v - w) \quad (15)$$

At the conformal infinity we have $v = w$, therefore $x' = x$, but, for $x \neq 0$, the coordinates v and w change. If $v = w = 0$, then, after the generic translation, $v' = w' \neq 0$. The coordinate description of the 2-sphere \mathcal{N}_s , which is the common part of \mathcal{J}^+ and \mathcal{J}^- changes. What is invariant is the set $\mathcal{J}^+ \cup \mathcal{J}^-$, and the fact that \mathcal{J}^+ and \mathcal{J}^- have a common 2-sphere.

5.3 Transitivity of $ISO(3, 1)$ on the conformal infinity

Let \mathcal{J} denote the conformal infinity, *minus the singular point* $I_0 = I_+ = I_-$. It is easy to see that action of $ISO(3, 1)$ on the is transitive. \mathcal{J} has the topology of a cylinder $\mathbb{R} \times S^2$. The group of translations acts along the \mathbb{R} , while $SO(3, 1)$ acts transitively on S^2 in a standard way - Lorentz transformations act on directions of light rays through the origin of the Minkowski space. It follows that any splitting of \mathcal{J} into \mathcal{J}^+ and \mathcal{J}^- is not translation invariant and not intrinsic. The article of Roger Penrose [24] is extremely unclear in this respect. Penrose mentions for instance that “There is another version of compactified Minkowski space in which the future boundary hypersurface is identified with the past”, and quotes his earlier paper [25], as well as the classic one by Kuiper [26], but he does not bother to define precisely what would be the alternative for the projective model. The same lack of clarity concerns the discussion in [20] and [19]. B.G. Schmidt, in an apparently mathematically precise paper [13] proves a Theorem stating that *The conformal boundary of Minkowski space is $\mathcal{J}^+ \cup \mathcal{J}^- \cup I^+ \cup I^- \cup I^0$* , without ever bothering to define the sets on the right hand side of his statement.

In [27, p. 178] Penrose writes:

“There is one property of \mathcal{R} , however, which seems undesirable when these ideas are applied to interacting fields, or curved space–times. This is the fact that the ‘future infinity’ turns out to have been identified with the ‘past infinity’ in the definition of \mathcal{R} . To avoid this feature it will be desirable effectively to ‘cut’ this manifold along the hypersurface \mathcal{J} and to consider instead the resulting manifold with boundary. This boundary consists essentially of two copies of \mathcal{J} , one in ‘future’ which will be called \mathcal{J}^+ and one in the ‘past’ to be called \mathcal{J}^- ...”

Nowhere a precise definition of \mathcal{J}^+ and \mathcal{J}^- is given. We are not told how the Poincaré group acts on these ‘boundaries’. Also the authors of recent papers like, for instance [28], when asked about the definition of \mathcal{J}^+ and \mathcal{J}^- for Minkowski space, refer to Penrose [27] or [29]. In fact Geroch does not define \mathcal{J}^+ and \mathcal{J}^- for the Minkowski space. He considers Schwarzschild space–time with the topology $S^2 \times \mathbb{R}^2$, proposes some coordinate-dependent constructions and does not really discuss global symmetries.

6 Light trapped at infinity

The aim of this section is to demonstrate that a light ray can be trapped in the conformal infinity and circulate there forever - unless disturbed by some quantum effect. It is well known (cf. e.g., [9] for a clear and self-contained exposition) that null geodesics are described by two-dimensional totally isotropic subspaces of $E^{4,2}$. Using the coordinates (x, v, w) as in Sec. 3.1, let x_0 be a fixed non-zero null vector in $E^{3,1}$, and let n_1 and n_2 be the vectors in $E^{4,2}$ defined by $n_1 = (x_0, 0, 0)$ and $n_2 = (0, 1, 1)$. Then the two-dimensional (real) plane spanned by n_1 and n_2 is totally isotropic - therefore it is describing a null geodesic in the compactified Minkowski space. A general vector in this plane is of the form $\alpha n_1 + \beta n_2 = (\alpha x_0, \beta, \beta)$, therefore it is completely contained in the conformal infinity that consists of null vectors (x, v, w) with $v = w$. We can completely parameterize our null geodesic by a parameter $\tau \in [0, \pi]$ by choosing the representatives of its points in the form

$$(\cos(\tau)x_0, \sin(\tau), \sin(\tau)). \quad (16)$$

For $\tau = 0$ the geodesic is on the 2-sphere \mathcal{N}_s , for $\tau = \pi/2$ it reaches the exceptional point $I^+ = I^- = I^0$, then it circulates further towards the 2-sphere \mathcal{N}_s . Notice that for $\tau = \pi$ we get the point $(-x_0, 0, 0)$ which projects

onto the same point of PN as $(x_0, 0, 0)$. Replacing x_0 by λx_0 , $\lambda \in \mathbb{R}$ does not change the plane spanned by n_1, n_2 , therefore in this way we get a family of null geodesics, all trapped in the conformal infinity. We can always choose a representative of x_0 of the form $(\mathbf{r}, 1)$, $\mathbf{r}^2 = 1$, so that we have a trapped null geodesic for every point of the unit sphere in \mathbb{R}^3 .

6.1 Conformal inversion

Consider the following linear map R of $E^{4,2}$: $R : (x, v, w) \mapsto (x, -v, w)$. It is clear that $R \in O(4, 2)$ (though not in $SO(4, 2)$). It is instructive to see that R implements the conformal inversion $x \mapsto x/x^2$ of the Minkowski space. To this end let x be a point in the Minkowski space M and let, writing x^2 for $q(x)$,

$$\tau(x) = \left(x, \frac{1}{2}(1 - x^2), -\frac{1}{2}(1 + x^2) \right)$$

be its image in $E^{4,2}$ as in Eq. (9).⁷ We apply the inversion R to obtain $(x, -\frac{1}{2}(1 - x^2), -\frac{1}{2}(1 + x^2))$ and represent it as an image of a new point x' . Therefore we should have

$$\left(x', -\frac{1}{2}(1 - x'^2), -\frac{1}{2}(1 + x'^2) \right) = \lambda \left(x, \frac{1}{2}(1 - x^2), -\frac{1}{2}(1 + x^2) \right). \quad (17)$$

Now, from $x' = \lambda x$ it follows that $x'^2 = \lambda^2 x^2$. Substituting this value of x'^2 into the two other equations and adding them we get $\lambda = 1/x^2$, therefore

$$x' = \frac{x}{x^2},$$

which is the well known conformal inversion in Minkowski space. The formula (17) becomes then an identity.⁸

Let us now apply the conformal inversion R to the light rays circulating at infinity, given by the formula (16). We obtain the family

$$(x_0, -\sin(\tau), \sin(\tau)) = -2 \sin(\tau) \left(x(\tau), \frac{1}{2}(1 - x(\tau)^2), -\frac{1}{2}(1 + x(\tau)^2) \right),$$

⁷This is the standard map, discussed in a general signature for instance in [23, p. 92].

⁸It is evident that this formula makes sense only when a length scale is chosen. This can be a Planck length, a cosmic scale length or some other length scale. The formula is singular on the light cone, but this apparent singularity is a coordinate effect.

where $x(\tau) = -\frac{1}{2} \cot(\tau)x_0$. This is nothing else but a family of light rays through the origin of the Minkowski space in the directions of null vectors x_0 . The parameter τ is, of course, not an affine parameter of these null geodesics.

6.2 The signature of the metric at infinity

Let H_1 be the affine hyperplane in $E^{4,2}$ parameterized by the coordinates (\mathbf{r}, t, v, w) , defined by the condition $t = 1$. Then H_1 is transversal with respect to the null cone \mathcal{N} , therefore, by Theorem 3 of [9] it induces the unique conformal structure on $\pi(H_1 \cap \mathcal{N})$. The intersection $H_1 \cap \mathcal{N}$ is described by the equation $\mathbf{r}^2 - 1 + v^2 - w^2 = 0$. Taking a trajectory there, by differentiation, we get for the tangent vector $(\dot{\mathbf{r}}, \dot{v}, \dot{w})$ the equation $\dot{\mathbf{r}} + \dot{v} - \dot{w} = 0$. Notice that at the points corresponding to the conformal infinity we have $v = w$. Taking a trajectory with $v = w = \text{const}$ we get a trajectory on the 2-sphere. The signature there is $(2, 0)$. On the other hand, taking a trajectory with \mathbf{r} constant we obtain a tangent vector of the form $(\mathbf{0}, 0, \dot{v}, \dot{w})$ - a null vector in $E^{4,2}$. It follows that the metric induced on conformal infinity is degenerate and has as its standard form $\text{diag}(1, 1, 0)$

6.3 A pictorial representation of the infinity

In order to get an idea about the manifold structure of the conformal infinity and to obtain its pictorial representation, it is convenient to use the formulas from Lemma 2. At the conformal boundary we have $v = w$, thus $v = \cos(\psi)$, and since $v^2 + \mathbf{x}^2 = 1$, we get $\mathbf{x}^2 = \sin^2(\psi)$. Furthermore, because (\mathbf{x}, t, v, w) and $(-\mathbf{x}, -t, -v, -w)$ describe the same point of $P\mathcal{N}$, it is enough to consider $\psi \in [0, \pi]$. The whole conformal infinity is then described by one equation:

$$\mathbf{x}^2 = \sin^2(\psi), \quad \psi \in [0, \pi],$$

where $(\mathbf{x}, \psi = 0)$ and $(\mathbf{x}, \psi = \pi)$ describe the same point. This is nothing else but a squeezed torus. Replacing the spheres S^2 by circles S^1 we get the graphic representation as shown in Fig. 6. Topology itself is represented by a double cone with two vertices identified, as in Fig. 3. This picture must not be confused with a similarly looking picture taken from [4, p. 178], which we reproduce here in Fig. 4.

6.4 The double covering

It is possible to repeat the constructions of Sects 3.1 and 4, but replacing the equivalence relation $Z \sim Z'$ by a stronger one: we identify two vectors Z and Z' in $E^{4,2}$ if and only if $Z' = \lambda Z$, $\lambda > 0$. The manifold resulting by taking the quotient of \mathcal{N} by this new equivalence relation will be denoted by $\hat{P}\mathcal{N}$. Instead of one map τ as in Eq. (9) we define now two maps:

$$\tau_+(x) = \left(x, \frac{1}{2}(1 - q(x)), -\frac{1}{2}(1 + q(x))\right). \quad (18)$$

$$\tau_-(x) = \left(x, -\frac{1}{2}(1 - q(x)), \frac{1}{2}(1 + q(x))\right). \quad (19)$$

Similarly we define

$$\mathcal{Z}_\pm = \{Z \in E^{4,2} : v - w = \pm 1\}, \quad (20)$$

and then show that

Lemma 3. *The image $\tau_\pm(E^{3,1})$ in $E^{4,2}$ coincides with the intersection $\mathcal{N} \cap \mathcal{Z}_\pm$ of \mathcal{N} with \mathcal{Z}_\pm .*

The manifold $\hat{P}\mathcal{N}$ contains now two copies of Minkowski space, we may call them M_+ and M_- , joined by a common boundary.

In Figure 2 the corresponding Penrose diagram is shown, this time we have two different 2-spheres S^+ and S^- . There are two copies of Minkowski space, M_+ and M_- , separated by the boundary. The horizontal lines at $\psi = +\pi$ and $\psi = -\pi$ should be identified. The corresponding pictorial representation of the infinity is shown in Figures 3, 6.

7 Geometry of oriented twistors

In this section we present a slightly modified version of the reasoning of Kopczyński and Woronowicz in [9, section III]⁹. In particular will take into account the orientation, and also we will change the notation a little bit by introducing the Hodge \star operator. Otherwise, in this section we will follow

⁹Our numbering conventions differ slightly from those used in [9]. We use Roman letters e, x, y, v, w , etc. to denote the elements of the algebra.




Figure 2: The second version of the “Penrose diagram” of Minkowski space.

the notation of the [9] - that may differ from the notation in other parts of this paper. To start with: as it will be explicitly shown below in section 7.2.2, twistors are spinors for the conformal group¹⁰. But, for our present purpose, in order to analyze the twistor geometry no knowledge of spinors is needed. We will make this section self-contained - to a large extent.

7.1 The exterior algebra $\bigwedge V$ and Hodge duality operator

Let V be a complex vector space of finite dimension n . We denote by $\bigwedge V = \bigoplus_{k=0}^n \bigwedge^k V$ the exterior algebra of V thought of as a consisting of antisym-

¹⁰In his Afterward to “Such Silver Currents. The Story of William and Lucy Clifford 1845-1929” [30, p. 182] Roger Penrose wrote: *Twistors may be regarded as spinors for six dimensions; yet they refer directly to the four dimensions of space-time.* In “The Road to Reality” [31, Ch. 33.4] Penrose writes *How do twistors fit in with all this? The shortest but hardly the most transparent way to describe a (Minkowski-space) twistor is to say that it is a reduced spinor (or half spinor) for $O(2, 4)$.*

metric tensors endowed with the wedge product¹¹:

$$v_1 \wedge \dots \wedge v_k = \sum_{\sigma} (-1)^{\sigma} v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(k)}.$$

Assume that V is endowed with a pseudo-hermitian form $(x|y)$ of signature (p, q) . The standard example is the space $\mathbb{C}^n = \mathbb{C}^p \oplus \mathbb{C}^q$ with

$$(x|y) = \sum_{i=1}^p x^i \bar{y}^i - \sum_{j=1}^q x^j \bar{y}^j.$$

We endow $\bigwedge V$ with a natural pseudo-hermitian form defined by:

$$(v_1 \wedge \dots \wedge v_k | w_1 \wedge \dots \wedge w_k) = \det((v_i | w_j)). \quad (21)$$

Remark 5. Notice that there exist, in the literature, two different conventions of defining the exterior product. While most authors seem to agree on the definition of the alternating operator:

$$\text{Alt}(v_1 \otimes \dots \otimes v_k) = \frac{1}{k!} \sum_{\sigma} (-1)^{\sigma} v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(k)},$$

the exterior product of a k -vector v and l -vector w can be defined by the formula:

$$v \wedge w = \left(\frac{(k+l)!}{k!l!} \right)^{\epsilon} \text{Alt}(v \otimes w),$$

where $\epsilon = 0$ or $\epsilon = 1$. We choose $\epsilon = 1$.

There are also two different convention of extending the scalar product from V to $\bigwedge V$. Some authors (especially physicists, when discussing the second quantization of Fermions) endow $\bigwedge V$ with the restriction of the natural scalar product defined on the tensor product. For k -vectors this gives $k!$ times our scalar product.

Given $x \in \bigwedge^p V$ we have the coordinate representation of x in a basis $\{e_i\}$ of V :

$$x = \frac{1}{p!} x^{i_1 \dots i_p} e_{i_1} \wedge \dots \wedge e_{i_p}.$$

¹¹For more information about exterior (Grassmann) algebras see e.g., [32, Ch. 5].

The wedge product is then given by the formula:

$$(x \wedge y)^{i_1 \dots i_{p+q}} = \frac{1}{p!q!} \delta_{j_1 \dots j_p j_{p+1} \dots j_{p+q}}^{i_1 \dots i_{p+q}} x^{j_1 \dots j_p} y^{j_{p+1} \dots j_{p+q}},$$

where $\delta_{c \dots d}^{a \dots b}$ is the (generalized) Kronecker delta symbol. We also have the coordinate representation:

$$(x|y) = \frac{1}{p!} G_{i_1 j_1} \dots G_{i_p j_p} x^{i_1 \dots i_p} \overline{y^{j_1 \dots j_p}}, \quad (22)$$

where $G_{ij} = \langle e_i, e_j \rangle$.

Let now $\{e_i\}$ be an orthonormal basis for V with $(e_i|e_i) = +1$ for $i = 1, \dots, p$, and $= -1$ for $i = p+1, \dots, p+q$, and let $e = e_1 \wedge \dots \wedge e_n$. Then $(e|e) = (-1)^q$. Let $e \in \bigwedge^n V$ be a unit n -vector. We call e an *orientation* of V . An orthonormal basis $\{e_i\}$ will be called *oriented* if $e_1 \wedge \dots \wedge e_n = e$. Any two oriented bases are then related by a unique transformation from the group $SU(r, s)$.

For each $x \in \bigwedge V$ let $C(x)$ be the linear operator on $\bigwedge V$ defined by

$$C(x)y = x \wedge y.$$

Clearly, for $x \in \bigwedge^k V$, we have $C(x) : \bigwedge^l V \rightarrow \bigwedge^{k+l} V$, and $v \mapsto C(v)$, $v \in V$ is a linear map from V to $L(\bigwedge V)$, with

$$C(v)C(w) + C(w)C(v) = 0, \quad (23)$$

for all $v, w \in V$. Notice that it follows from the definition that $C(x \wedge y) = C(x)C(y)$.

Let $C(x)^*$ be the Hermitian adjoint of $C(x)$, defined by $(C(x)^*y|z) = (y|C(x)z)$, $y, z \in \bigwedge V$. Then, for $x \in \bigwedge^k V$, $C(x)^* : \bigwedge^l V \rightarrow \bigwedge^{l-k} V$, the map $x \mapsto C(x)^*$ is anti-linear, and for $v, w \in V$ we have the anti-commutation relations:

$$C(v)C(w)^* + C(w)^*C(v) = (v|w). \quad (24)$$

Notice that for all $x, y \in \bigwedge V$ we have $C(x \wedge y)^* = C(y)^*C(x)^*$.

Remark 6. *The anti-commutation relations (23,24) are known as CAR - canonical commutation relations - in our case finite-dimensional and generalized for the case of an indefinite scalar product. If we define $\phi(v) =$*

$2^{-\frac{1}{2}}(C(v) + C(v)^*)$, then the real linear map $v \mapsto \phi(v)$ is a Clifford map for V considered as a $2n$ -dimensional real vector space endowed with the scalar product $\Re((v|w))$ - cf. [33]¹²

Assuming V oriented with an orientation e , we define the Hodge operator $\star : \bigwedge^k V \rightarrow \bigwedge^{n-k} V$ as an antilinear map $\star : x \mapsto \star x$ uniquely defined by the formula

$$x \wedge \star y = (x|y)e, \quad x, y \in \bigwedge^k V. \quad (25)$$

It is easy to see that an equivalent definition of the Hodge \star operator is given by:

$$\star x = C(x)^* e.$$

It easily follows from the definition that for $x \in \bigwedge^k V, y \in \bigwedge^{n-k} V$ we have:

$$(x|\star y) = (-1)^{k(n-k)}(y|\star x). \quad (26)$$

A little bit more effort¹³ is required to check that we have

$$\star \star x = (-1)^{k(n-k)+q} x, \quad \forall x \in \bigwedge^k V.$$

Remark 7. A k -vector $x \neq 0$ is called **decomposable** if x is of the form $x = x_1 \wedge \dots \wedge x_k$ for $x_1, \dots, x_k \in V$. If x is decomposable, then also $\star x$ is decomposable. Moreover the $(n-k)$ -dimensional subspace corresponding to $\star x$ is the orthogonal complement of the subspace corresponding to x - cf. [42, Exercise 8, p. 62].

Another important property involving creation and annihilation operators to the Hodge star operator is [10, eq. 139] is¹⁴

$$\star C(x)^* \star^{-1} = C(x)(-1)^{d(x)\hat{N}},$$

where $d(x)$ is the grade of x ($d(x) = k$ for $x \in \bigwedge^k V$) and \hat{N} is the number operator - $\hat{N}y = ly$ for $y \in \bigwedge^l V$.

¹²For a complex number $z = \alpha + i\beta$ we denote $\Re(z) = \alpha, \Im(z) = \beta$.

¹³Cf. e.g., [34, p. 167], [35, p. 118]

¹⁴While only positive definite scalar product is discussed in [10], this particular property can be easily seen to hold also for pseudo-Hermitian spaces.

We define a *bilinear* form

$$\langle x, y \rangle = (x | \star y), \quad x, y \in \bigwedge V.$$

Notice that the following formulas hold:

$$\begin{aligned} \langle x, y \rangle &= (-1)^q \langle y, x \rangle \\ \overline{\langle x, y \rangle} &= (-1)^{k(n-k)+q} \langle x, y \rangle \end{aligned}$$

In an orthonormal basis e_i such that $e = e_1 \wedge \dots \wedge e_n$ we have the explicit expression for the star operator for $x \in \bigwedge^p V$:

$$(\star x)^{i_{p+1} \dots i_n} = \frac{1}{p!} G_{i_1 j_1} \dots G_{i_p j_p} \epsilon^{i_1 \dots i_p i_{p+1} \dots i_n} \overline{x^{j_1 \dots j_p}}. \quad (27)$$

7.2 The case of signature (2, 2)

In this section we specialize to the case of the signature (2, 2) that is relevant for our purposes, and has been studied in [9].

Let G be the diagonal matrix $G = \text{diag}(+1, +1, -1, -1)$. Let $H_{2,2}$ be a four-dimensional complex vector space endowed with a pseudo-Hermitian form $(\cdot | \cdot)$ of signature (2,2). A basis e_i of $H_{2,2}$ is said to be orthonormal if $(e_i | e_j) = G_{ij}$. Any two orthonormal bases are related by a transformation from the group $U(2, 2)$. We fix an orientation $e \in \bigwedge^4 H_{2,2}$ and define the Hodge \star duality operator as in previous subsection). Notice that $\bigwedge^2 H_{2,2}$ we have $\star^2 = 1$. Let $\Re \bigwedge^2 H_{2,2}$ be the space of self-dual bivectors:

$$\Re \bigwedge^2 H_{2,2} = \{x \in \bigwedge^2 H_{2,2} : x = \star x\}.$$

Then $\Re \bigwedge^2 H_{2,2}$ is a six-dimensional real vector space, and the real-bilinear form $\langle x, y \rangle$ is real-valued and symmetric on $\Re \bigwedge^2 H_{2,2}$. It can be easily seen (Cf. [9, Theorem 7]) that $\Re \bigwedge^2 H_{2,2}$ equipped with the scalar product $\langle x, y \rangle$ is of signature (4, 2). It follows that all the constructions of section 6.4 apply and in the following we will use the notation of this section. In particular we will the identification $E^{4,2} = \Re \bigwedge^2 H_{2,2}$.

In a complex vector space the concept of an orientation of a subspace is not well defined. In our case, however we can define what is meant by an oriented two-dimensional subspace. Given a k -dimensional subspace S

we can associate with it a simple (i.d. decomposable) nonzero k -vector x , unique up to a non-zero complex factor. For $\lambda \neq 0$ x and λx define the same subspace. For $k = 2$ we can restrict the freedom of choice by demanding that x should be self-dual: $\star x = x$. This restricts the freedom of choice to λ real - that is either positive or negative. By an “oriented two-space” we will thus mean an equivalence class of simple self-dual bivectors, where x and y define the same oriented subspace if and only if $y = \lambda x$, $\lambda > 0$.

Consider now the Grassmann manifold of oriented *totally isotropic* (complex) subspaces of $H_{2,2}$. We can repeat now, slightly modified, argument of [9].¹⁵

Theorem 1. *There is a one-to-one correspondence between the elements of $\hat{P}\mathcal{N}$ (the double covering of the compactified Minkowski space), and the oriented isotropic subspaces of $H_{2,2}$.*

Proof. If $p \in \hat{P}\mathcal{N}$, then there exists a unique up to a multiplication by a positive constant, non-zero element x of $E^{4,2}$ in the equivalence class of p . Since x is a null vector of $E^{4,2}$, and since, as a bivector, it is self-dual, it follows that $x \wedge x = x \wedge \star x = (x|x)e = (x|\star x)e = \langle x, x \rangle e = 0$. Therefore x is decomposable and it represents a two-dimensional subspace $S(q)$. Now, since x is self dual, $x = \star x$, it follows from the Remark 7 that $S(q)$ is orthogonal to itself, and thus totally isotropic as a subspace of $H_{2,2}$. Conversely, let x be a self-dual bivector representing an oriented totally isotropic subspace S . Then $(x|x) = 0$ (since the subspace is totally isotropic), and, since $\star x = x$, we have $\langle x, x \rangle = 0$, thus x is an isotropic vector of $E^{4,2}$, and therefore determines $p \in \hat{P}\mathcal{N}$. \square

7.2.1 Relation to the $U(2)$ compactification

In section 2 the points of \tilde{M} have been described by unitary operators $U \in U(2)$, while in this Section by rays in the space of self-dual null bivectors in $E^{4,2}$. It may be of interest to derive an explicit formula connecting these two descriptions.

Let us equip $H_{2,2}$ with an orthonormal basis e_1, e_2, e_3, e_4 and orientation $e = e_1 \wedge \dots \wedge e_4$. Then $H_{2,2}$ can be decomposed into $H_{2,2} \approx \mathbb{C}^2 \oplus \mathbb{C}^2$, and every vector $x \in H_{2,2}$ can be written as

$$x = \begin{pmatrix} u \\ v \end{pmatrix}, \quad u, v \in \mathbb{C}^2.$$

¹⁵For an additional information related to this subject, see also [6, 36].

It is easy to see that there is a bijection between unitary matrices U in \mathbb{C}^2 and maximal totally isotropic subspaces in $H_{2,2}$: Every maximal totally isotropic subspace \mathcal{W} of $H_{2,2}$ is of the form:

$$\mathcal{W} = \left\{ \begin{pmatrix} Uv \\ v \end{pmatrix} : v \in \mathbb{C}^2 \right\},$$

where U is uniquely determined by \mathcal{W} . Conversely, given unitary U the above formula defines a 2-dimensional maximal totally isotropic subspace \mathcal{W} . For our purposes it will be convenient to write the unitary operators as cU , where c is in $U(1)$, (i.e., $\{c \in \mathbb{C} : |c| = 1\}$), and U is in $SU(2)$. To each $(c, U) \in U(1) \times SU(2)$ we associate a maximal totally isotropic subspace $\mathcal{W}(c, U)$ defined by

$$\mathcal{W}(c, U) = \left\{ \begin{pmatrix} Uv \\ cv \end{pmatrix} : v \in \mathbb{C}^2 \right\}.$$

Till now we still have a redundancy, since (c, U) and $(-c, -U)$ define the same subspace. However, this redundancy will soon disappear when we will move from subspaces to oriented bivectors. In order to do this select two basis vectors in \mathbb{C}^2 :

$$v_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix},$$

and let $f_i(c, U) \in H_{2,2}$, $i = 1, 2$ be defined by

$$f_i = \begin{pmatrix} Uv_i \\ cv_i \end{pmatrix}.$$

Every matrix $U \in SU(2)$ can be uniquely written in the form

$$U = \begin{pmatrix} \bar{\alpha} & \beta \\ -\bar{\beta} & \alpha \end{pmatrix}, \quad |\alpha|^2 + |\beta|^2 = 1.$$

Our vectors f_i can then be written in components as follows:

$$f_1 = \begin{pmatrix} \bar{\alpha} \\ -\bar{\beta} \\ c \\ 0 \end{pmatrix}, \quad f_2 = \begin{pmatrix} \beta \\ \alpha \\ 0 \\ c \end{pmatrix},$$

or $f_1 = \alpha e_1 - \bar{\beta} e_2 + c e_3$ and $f_2 = \beta e_1 + \bar{\alpha} e_2 + c e_4$. To the pair (c, U) we associate the bivector $f_1 \wedge f_2$, easily calculated to be

$$f_1 \wedge f_2 = e_{12} - c\beta e_{13} + c\bar{\alpha} e_{14} - c\alpha e_{23} - c\bar{\beta} e_{24} + c^2 e_{34},$$

where $e_{ij} = e_i \wedge e_j$. It follows by the very construction that $f_1 \wedge f_2$ is a null vector in $\bigwedge^2 H_{2,2}$, what can be easily checked, but it is not, in general, self-dual: $\star(f_1 \wedge f_2) \neq f_1 \wedge f_2$. Therefore let us consider bivector f defined by the formula:

$$f = \frac{1}{\sqrt{2}} (f_1 \wedge f_2 + \star(f_1 \wedge f_2)).$$

Now $f(c, U)$ is both null and self-dual.

From the explicit formulas (22), (27) we easily find the following properties of the basis vectors e_i , $i = 1, \dots, 4$:

$$1 = (e_{12}|e_{12}) = (e_{34}|e_{34}) = -(e_{13}|e_{13}) = -(e_{14}|e_{14}) = -(e_{23}|e_{23}) = -(e_{24}|e_{24}),$$

and $\star e_{12} = e_{34}$, $\star e_{13} = e_{24}$, $\star e_{14} = -e_{23}$, $\star e_{23} = -e_{14}$, $\star e_{24} = e_{13}$, $\star e_{34} = e_{12}$. Define the following six 4×4 antisymmetric matrices $\Sigma_i = (\Sigma_i^{AB})$:¹⁶

$$\Sigma_1 = \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \\ i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix}$$

$$\Sigma_2 = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

$$\Sigma_3 = \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix}$$

$$\Sigma_4 = \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \end{pmatrix}$$

¹⁶These matrices have been constructed using the fact that $Cl_{4,2} = Cl_{3,1} \otimes Cl_{1,2} = (Cl_{2,0} \otimes Cl_{1,1}) \otimes Cl_{1,1}$, constructing this way 8×8 real matrix generators, then finding a unique (up to scale) metric matrix invariant under the Clifford group - of signature $(4,4)$, and also a unique invariant complex structure in \mathbb{R}^8 , the expressing the generators as complex 4×4 matrices, and renumbering them generators.

$$\Sigma_5 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$$

$$\Sigma_6 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}$$

Lemma 4. *The following identities hold:*

$$\overline{\Sigma_\alpha^{ij}} = \frac{1}{2} \epsilon^{ijkl} G_{km} G_{ln} \Sigma_\alpha^{mn}, \quad (28)$$

where $\alpha = 1, \dots, 6$, $i, j, k, l, m, n = 1, \dots, 4$.

Proof. Easily follows by a direct calculation.¹⁷ \square

It follows from the Eq. (28) that if we define bivectors E_1, \dots, E_6 by the formula

$$E_\alpha = \frac{1}{2\sqrt{2}} \Sigma_\alpha^{ij} e_i \wedge e_j,$$

then

$$\star E_\alpha = E_\alpha.$$

Moreover, one can verify that we have

$$\langle E_\alpha, E_\beta \rangle = Q_{\alpha\beta},$$

where

$$Q = \text{diag} (1, 1, 1, -1, 1, -1). \quad (29)$$

Explicitly we have:

$$\begin{aligned} E_1 &= \frac{i}{\sqrt{2}}(e_{13} - e_{24}), & e_{12} &= \frac{-1}{\sqrt{2}}(E_6 - iE_4) \\ E_2 &= \frac{1}{\sqrt{2}}(e_{13} + e_{24}), & e_{13} &= \frac{1}{\sqrt{2}}(E_2 - iE_1) \\ E_3 &= \frac{-i}{\sqrt{2}}(e_{14} + e_{23}), & e_{14} &= \frac{-1}{\sqrt{2}}(E_5 - iE_3) \\ E_4 &= \frac{-i}{\sqrt{2}}(e_{12} - e_{34}), & e_{23} &= \frac{1}{\sqrt{2}}(E_5 + iE_3) \\ E_5 &= \frac{-1}{\sqrt{2}}(e_{14} - e_{23}), & e_{24} &= \frac{1}{\sqrt{2}}(E_2 + iE_1) \\ E_6 &= \frac{-1}{\sqrt{2}}(e_{12} + e_{34}), & e_{34} &= \frac{-1}{\sqrt{2}}(E_6 + iE_4) \end{aligned}$$

¹⁷These and other calculations in this paper have been aided by several different computer algebra systems.

Then, the calculation gives the following result:

$$f = -\sqrt{2}\Re(c) (\Im(\beta)E_1 + \Re(\beta)E_2 - \Im(\alpha)E_3 - \Im(c)E_4 + \Re(\alpha)E_5 + \Re(c)E_6)$$

Evidently there is a problem with this definition for $\Re(c) = 0$. But we are free to choose the scale factor in our definition, therefore we *define*:

$$f(c, U) \stackrel{df}{=} \Im(\beta)E_1 + \Re(\beta)E_2 - \Im(\alpha)E_3 - \Im(c)E_4 + \Re(\alpha)E_5 + \Re(c)E_6. \quad (30)$$

It is easy to see that the formula above provides an embedding of $U(1) \times SU(2)$ into the isotropic cone \mathcal{N} of $E(4, 2)$ that is transversal to the generator lines of \mathcal{N} , and induces a diffeomorphism from $U(1) \times SU(2)$ onto $\hat{P}\mathcal{N}$. Notice that we have $f(-c, -U) = -f(c, U)$, thus replacing $c \mapsto -c$, $U \mapsto -U$ changes the orientation of the corresponding isotropic subspace.

Let us now return to the formula (5) of sec. 2 that provides the embedding ψ of M into $U(2)$ via the Cayley transform. We rewrite it in the form

$$\psi(x) = cU', \quad c \in U(1), \quad U' \in SU(2)$$

where

$$c = \frac{-(1 + q(x) + 2ix^0)}{\sqrt{(1 + q(x))^2 + 4x^{02}}},$$

$$U' = \frac{1}{\sqrt{(1 + q(x))^2 + 4x^{02}}} \begin{bmatrix} 1 - q(x) + 2ix^3 & 2(ix^1 + x^2) \\ 2(ix^1 - x^2) & 1 - q(x) - 2ix^3 \end{bmatrix}.$$

Applying the formula (30) we obtain

$$f(c, U') = \lambda (x^1 E_1 + x^2 E_2 + x^3 E^3 + x^0 E_4) + \lambda \left(\frac{1}{2}(1 - q(x))E_5 - \frac{1}{2}(1 + q(x))E_6 \right),$$

where

$$\lambda = \frac{2\sqrt{2}}{(1 + q(x))^2 + 4x^{02}} > 0.$$

This is the same map as the one given by Eq. (9).

7.2.2 From self–dual bivectors to the Clifford algebra and conformal spinors

In Chapter 1.5.5.1 of [23] Pierre Anglès generalizes earlier results of Deheuvels and shows how to embed the projective null cone of $E^{p,q}$ into the space of spinors of the Clifford algebra of this space. It is instructive to see how this method works in our case, yet in order to this we must first explicitly identify the space of spinors for our version of $E^{4,2}$ realized as self–dual bivectors in $H_{2,2}$.

Proposition 1. *Define the following six complex matrices*

$$\Gamma_{\alpha k}^i = \Sigma_{\alpha}^{ij} Q_{jk}, \quad (\alpha = 1, \dots, 6; i, j, k = 1, \dots, 4) \quad (31)$$

and let Γ_{α} be the antilinear operators on $H_{2,2}$ defined by the formula:

$$(\Gamma_{\alpha} f)^i = (\Gamma_{\alpha})_j^i \overline{f^j}, \quad f = (f^i) \in H_{2,2}.$$

Then the antilinear operators Γ_{α} satisfy the following anti–commutation relations of the Clifford algebra of $E^{4,2}$:

$$\Gamma_{\alpha} \circ \Gamma_{\beta} + \Gamma_{\beta} \circ \Gamma_{\alpha} = 2 Q_{\alpha\beta}.$$

The space $H_{2,2}$ considered as an 8–dimensional real vectors space carries this way an irreducible representation of the Clifford algebra $Cl_{4,2}$. The Hermitian conjugation in $H_{2,2}$ coincides with the main anti–automorphism of $Cl_{4,2}$. The space $H_{2,2}$ considered as a 4–dimensional complex vector space carries a faithful irreducible representation of the even Clifford algebra $Cl_{4,2}^+$.

Proof. The formulas (31) follow easily by a direct calculation. The first part of the Proposition follows then from the known fact that the Clifford algebra $Cl_{4,2}$ is isomorphic to the algebra $Mat(8, \mathbb{R})$, while the even Clifford $Cl_{4,2}^+$ is known to be isomorphic to $Mat(4, \mathbb{C})$ (cf. e.g., [23, Table 1.1, p. 28]). Moreover, also by the direct calculation we have

$$(\Gamma_{\alpha} \circ \Gamma_{\beta})^* = \Gamma_{\beta} \circ \Gamma_{\alpha},$$

which proves the statement about the main automorphism. \square

Remark 8. *It follows from the above Proposition that $H_{2,2}$ is a spinor space for the Clifford algebra of its self–dual bivectors.*

In [23, Ch. 1.5.5.1, p. 44] Pierre Anglès discusses a general method of embedding a projective quadric into the manifold of totally isotropic subspaces of a spinor space for the even Clifford algebra. Let us apply this method to our case adding at the same time a new element to this method. The original method can be described as follows: Consider $E^{p,q}$ as a vector subspace of its Clifford algebra $Cl_{p,q}$. Let S be a spinor space for $Cl_{p,q}^+$ endowed with its associated scalar product. For each non-zero isotropic vector $x \in E^{p,q}$ find another isotropic vector y such that $2\langle x, y \rangle = 1$. Then yx is an idempotent in $Cl_{p,q}^+$, and its kernel $S(x)$ is a totally isotropic subspace of S that depends only on x and not on y . One disadvantage of this procedure in applications is that we are not being given a procedure for selecting y for each given x . This can be, however, in our case, easily improved.

Let us first describe the philosophy behind our procedure.¹⁸ The set D of maximal positive subspaces of $H_{2,2}$ is a complex symmetric domain for $U(2,2)$, $D = U(2,2)/(U(2) \times U(2))$, and the manifold of maximal totally isotropic subspaces is its Shilov's boundary \hat{D} . There is a one-to-one correspondence between maximal subspaces and Hermitian unitary operators J in $H_{2,2}$ with the property that the scalar product $(x|Jy)$ is positive definite on $H_{2,2}$. If J is such an operator, then the associated maximal positive subspace is given by $\{z \in H_{2,2} : Jz = z\}$. Every such J is, in particular, an element of $SU(2,2)$, therefore it acts on its Shilov's boundary \hat{D} . Acting on a given element of \hat{D} , it produces another element, its "J-antipode". We will take for J the operator described by the matrix G . It is then easy to see that in terms of isotropic vectors $E^{4,2}$ the corresponding action consists of flipping the signs of two coordinates: $(\mathbf{x}, t, x^5, x^6) \mapsto (\mathbf{x}, -t, x^5, -x^6)$. In other words - it corresponds to the action of the matrix Q - cf. (29).

Proposition 2. *Let x be a point in M , $x = (x^0, \mathbf{x})$, let $X = \tau(x)$ be its image in $E^{4,2}$, as in Eq. (9), and let $Y' = QX$ be its antipode. Let $Y = Y'/(2\langle X, Y' \rangle)$, so that $2\langle X, Y \rangle = 1$. Let $\hat{X} = X^1\Gamma_1 + \dots + X^6\Gamma_6$ be the image of X in $Cl_{4,2}$, and similarly for $\hat{Y} = Y^1\Gamma_1 + \dots + Y^6\Gamma_6$. Then $P = \hat{Y} \circ \hat{X}$ is an idempotent in $L(H_{2,2})$ whose kernel is a maximal totally isotropic subspace of $H_{2,2}$ consisting of vectors of the form (U^v) , where U is the unitary matrix given by Eq. (5).*

Proof. The proof follows by a straightforward though lengthy direct calculation. □

¹⁸For more information cf. [38] and references therein.

8 Flat conformal structures

While the present paper concentrates on the Minkowski space, the results apply also to tangent space structures in more general case - they may also apply to conformally flat manifolds. In this section we will introduce the main concepts needed for such an extension and show that the embedding τ given by Eq. 9 of section 3.1 can be understood geometrically by the *conformal development* with respect to the normal Cartan connection.

8.1 The bundle $P^2(M)$

Let M be a smooth n -dimensional manifold. Two maps from open neighborhoods of the origin $0 \in \mathbb{R}^n$ to M define the same 2-jet at 0 if and only if their partial derivatives up to the second order coincide. The 2-jet determined by such a map e is denoted $j_0^2(e)$. If e is a diffeomorphism, then $j_0^2(e)$ is called a second order frame at the point $p = e(0)$. The set of all second-order frames is denoted by $P^2(M)$.¹⁹

Let (x^μ) be a local chart of M , and let (t^a) be the standard coordinates on \mathbb{R}^n . Given $j_0^2(e)$ such that p is in the domain of the chart, a set of coordinates of $j_0^2(e)$ is defined by:

$$\begin{cases} e^\mu & \doteq x^\mu(p) \\ e^\mu_a & \doteq \left. \frac{\partial(x \circ e)^\mu}{\partial t^a} \right|_{t=0} \\ e^\mu_{ab} & \doteq \left. \frac{\partial^2(x \circ e)^\mu}{\partial t^a \partial t^b} \right|_{t=0} \end{cases}$$

If (x^μ) is replaced by $(x^{\mu'})$, the coordinates of $j_0^2(e)$ change:

$$\begin{cases} e^{\mu'} & = x^{\mu'}(p) \\ e^{\mu'}_a & = \frac{\partial x^{\mu'}}{\partial x^\mu}(p) e^\mu_a \\ e^{\mu'}_{ab} & = \frac{\partial x^{\mu'}}{\partial x^\mu}(p) e^\mu_{ab} + \frac{\partial^2 x^{\mu'}}{\partial x^\mu \partial x^\nu}(p) e^\mu_a e^\nu_b \end{cases}$$

It follows that e^μ_a may be considered as an ordinary (i.e., first order) frame at p . A natural projection $P^2(M) \rightarrow P^1(M)$ exists, and is given by $j_0^2(e) \mapsto j_0^1(e)$ or, in coordinates, by $(e^\mu, e^\mu_a, e^\mu_{ab}) \mapsto (e^\mu, e^\mu_a)$. A simple interpretation can be given to e^μ_{ab} . First notice that the matrix e^μ_a is always invertible. Let e^a_μ denote the inverse matrix, so that we have $e^\mu_a e^a_\nu = \delta^\mu_\nu$ and $e^a_\mu e^\mu_b = \delta^a_b$. Define ‘‘connection coordinates of e ’’ by

$$e^\mu_{\rho\sigma} \doteq -e^r_\rho e^s_\sigma e^\mu_{r\sigma}.$$

¹⁹For a somewhat different version cf. also [23, pp. 138-152]

It follows from the transformation properties of the coordinates of e above that $e^\mu_{\rho\sigma}$ transform as connection coefficients at p . Therefore each section of $P^2(M)$ determines a pair: a section of $P^1(M)$ (i.e., a frame) and a torsion-free affine connection on M , the correspondence being bijective. In particular, if $P^1(M)$ is reduced to the orthogonal or pseudo-orthogonal group, the Hilbert-Palatini principle for General Relativity can be considered as a functional on the space of sections of $P^2(M)$. Also notice that the diffeomorphisms group of M acts on $P^2(M)$ and on the space of its sections in a natural way. If e is a map from an open neighborhood of the origin $0 \in \mathbb{R}^n$ to M , and if $\phi : M \rightarrow M$ is a local diffeomorphism defined at $p = e(0)$, then $\phi \circ e$ is another map from an open neighborhood of the origin $0 \in \mathbb{R}^n$ to M . If e_1 and e_2 define the same second order frame: $j_0^2(e_1) = j_0^2(e_2)$, then the composed maps define the same second order frame as well: $j_0^2(\phi \circ e_1) = j_0^2(\phi \circ e_2)$.

8.2 The structure group $G^2(n)$

Let $G^2(n)$ denote the set of all second-order frames at $0 \in \mathbb{R}^n$. $G^2(n)$ is a group with the group multiplication law given by

$$j_0^2(h)j_0^2(k) \doteq j_0^2(h \circ k).$$

The group $G^2(n)$ acts on $P^2(M)$ from the right

$$j_0^2(e)j_0^2(h) \doteq j_0^2(e \circ h).$$

Corresponding to the canonical coordinates in \mathbb{R}^n , there are natural coordinates in $G^2(n)$: (h^a_b, h^a_{bc}) , and each $j_0^2(h)$ can be uniquely represented by the map $\mathbb{R}^n \rightarrow \mathbb{R}^n$ given by

$$t^a \mapsto h^a_r t^r + \frac{1}{2} h^a_{rs} t^r t^s.$$

In terms of natural coordinates the group composition law in $G^2(n)$ can be written as

$$(h^a_b, h^a_{bc})(k^a_b, k^a_{bc}) = (h^a_r k^r_b, h^a_{rs} k^r_b k^s_c + h^a_r k^r_{bc})$$

While the group $G^2(n)$ acts on $P^2(M)$ from the right, and $P^2(M)$ is a principal bundle over M with $G^2(n)$ as its structure group, the group $Diff(M)$ of diffeomorphisms of M acts on $P^2(M)$ from the left, by fibre preserving

transformations, commuting with the right action of $G^2(n)$ - thus as an automorphism group of $P^2(M)$. An affine connection can be considered as a section of a bundle associated to $P^2(M)$ via an appropriate representation of $G^2(n)$ by affine transformations.

8.3 Reduction of $P^2(M)$ induced by a conformal structure

Let now M be an orientable and oriented n -dimensional differentiable manifold. Let $GL_+(n)$ be the group of $n \times n$ real matrices of positive determinant. We denote by TM the tangent bundle of M , and by F_+ the $GL_+(n)$ principal bundle of oriented linear frames of M . We denote by Λ_+^n the bundle of oriented non-vanishing n -vectors. Λ_+^n is, in a natural way, a principal \mathbb{R}_+ bundle. Given a real number w , let V^w be the bundle associated to F_+ via the representation ρ_w of $GL_+(n)$ on \mathbb{R} defined by

$$\rho_w : GL_+(n) \ni A \mapsto \det(A)^{-w} \in \mathbb{R}.$$

Since any oriented frame e defines an oriented n -vector $e_1 \wedge \dots \wedge e_n$, it follows that V^w can be also considered as the bundle associated to Λ_+^n via the representation $\mathbb{R}_+ \ni x \mapsto x^w \mathbb{R}$.

Cross-sections of V^w are called densities of weight w . In what follows we will use the “hat” symbol $\hat{}$ to distinguish densities from tensorial objects of weight $w = 0$. If $e = \{e_i, i = 1, \dots, n\}$ is a frame at p , and if $\hat{\phi}$ is an element in the fibre V_p^w , then we denote by $\hat{\phi}[e]$ the real number representing $\hat{\phi}$ with respect to the frame e . We write $\hat{\phi} > 0$ if $\hat{\phi}[e] > 0$ for some (and thus for every) oriented frame. It follows from the very definition of the associated bundle that if $A \in GL_+(n)$, then $\hat{\phi}[eA] = \det[A]^w \hat{\phi}[e]$.

Let r, s be a pair of real numbers, and let $\hat{\phi}, \hat{\psi}$ be positive densities of weight $w = r$ and $w = s$ respectively. Then $(\hat{\phi}\hat{\psi})[e] = \hat{\phi}[e]\hat{\psi}[e]$ defines a density of weight $w = r + s$, while $\hat{\phi}^s[e] = \hat{\phi}[e]^s$ defines a density of weight $w = rs$.

Let $x^\mu, \mu = 1, \dots, n$ be a local coordinate system on M , and let ∂_μ be the basis made of vectors tangent to the coordinate lines. Then a cross-section $\hat{\phi}$ of V^w is represented by a real-valued function $\hat{\phi}(x)$. When the local coordinate system changes to another one, $x^{\mu'}$, then the coordinate bases changes accordingly:

$$\partial_{\mu'} = \frac{\partial x^\mu}{\partial x^{\mu'}} \partial_\mu,$$

and the corresponding numerical representation of $\hat{\phi}$ changes as follows:

$$\hat{\phi}'(x') = \left| \frac{\partial x'}{\partial x} \right|^{-w} \hat{\phi}(x),$$

where $\left| \frac{\partial x'}{\partial x} \right|$ is the Jacobian of the coordinate transformation.

By taking tensor products of tensor bundles with the line bundle V^w we can define, in an obvious way, tensor densities of weight w .

Although much of what will follow is true in a general case of an arbitrary (pseudo-)Riemannian manifold, we will assume in the following that we are dealing with the signature $(n-1, 1)$, that our manifold M is oriented and time-oriented, and that all our local coordinate systems have positive orientation and time-orientation.

Let $\eta_{ij} = \text{diag}(1, \dots, 1, -1, \dots, -1)$, (signature (p, q)) and let $O(\eta)$ be the subgroup of $GL(n)$ consisting of matrices $\Lambda = (\Lambda^i_j) \in GL(n)$ such that $\Lambda^t \eta \Lambda = \eta$, $\det \Lambda = 1$, and let $SO_0(\eta)$ be the connected component of the identity in $O(\eta)$. By a (pseudo-) Riemannian structure on M we will mean a reduction of the $GL(n)$ principal bundle of the linear frames of M to $SO_0(\eta)$.

There are several equivalent ways of defining a conformal structure on M . Probably the most intuitive way is to define it as “a Riemannian metric up to a scale”. Let g and \tilde{g} be two metrics of M . Then g and \tilde{g} are said to be conformally related if there exists a positive function $\hat{\phi}$ on M such that $\tilde{g}(p) = \hat{\phi}(p)g(p)$ for all $p \in M$. Being “conformally related” is, in fact, an equivalence relation, so that we can define a conformal structure on M as the equivalence class consisting of conformally related metrics.

Let C be a conformal structure on M . For any $g \in C$, given a local coordinate system x^μ , we can define $|g|$ to be the absolute value of the determinant $\det g_{\mu\nu}$, where $g_{\mu\nu} = g(\partial_\mu, \partial_\nu)$. Then from the transformation law:

$$g_{\mu'\nu'} = \frac{\partial x^\mu}{\partial x^{\mu'}} \frac{\partial x^\nu}{\partial x^{\nu'}} g_{\mu\nu}$$

we find that

$$|g'| = \left| \frac{\partial x}{\partial x'} \right|^2 |g|,$$

so that $|g|$ is a scalar density of weight -2 . Let us define

$$\gamma_{\mu\nu} = \frac{g_{\mu\nu}}{|g|^{1/n}}.$$

Then $\det \gamma_{\mu\nu} = -1$, $\gamma_{\mu\nu}$ is a symmetric tensor density of weight $-2/n$, and $\gamma_{\mu\nu}$ is independent of the choice of the representative $g_{\mu\nu}$ in the conformal class C . In other words: a conformal structure is uniquely characterized by a symmetric tensor density of weight $-2/n$, and signature (p, q) .

Let $\mathcal{T}M$ be the vector bundle of vector densities of weight $w = 1/n$. Then, for any two vectors $\hat{u}, \hat{v} \in \mathcal{T}_p M$ the number $(\hat{u}, \hat{v}) = \gamma_{\mu\nu} \hat{u}^\mu \hat{v}^\nu$ is independent of the local coordinate system at p - it defines a bilinear form of signature (p, q) on $\mathcal{T}M$. This bilinear form characterizes uniquely the conformal structure C .

Let a conformal structure C be given on M . A general torsion-free affine connection which preserves C is of the form

$$\Gamma_{\beta\gamma}^\alpha = \hat{\Gamma}_{\beta\gamma}^\alpha + (\delta_\beta^\alpha p_\gamma + \delta_\gamma^\alpha p_\beta - \gamma_{\beta\gamma} \gamma^{\alpha\rho} p_\rho),$$

where

$$\hat{\Gamma}_{\beta\gamma}^\alpha = \frac{1}{2} (\partial_\beta \gamma_{\gamma\rho} + \partial_\gamma \gamma_{\beta\rho} - \partial_\rho \gamma_{\beta\gamma}),$$

and $\gamma^{\mu\nu}$ is the inverse matrix of $\gamma_{\mu\nu}$. Therefore $P^2(M)$ can be reduced to $P_C^2(M)$ defined as consisting of second-order frames e such that (e^μ_a) are conformal frames and $e^\mu_{\rho\sigma}$ are the coefficients of conformal connections. It is easy to see that the structure group H of $P_C^2(M)$ is a subgroup of $G^2(n)$ consisting of pairs (h^a_b, h^a_{bc}) , with $h^a_b \in CO_0(\eta)$, and

$$h^a_{bc} = h^a_r (\delta_b^r v_c + \delta_c^r v_b - \eta_{bc} \eta^{rs} v_s),$$

where $CO_0(\eta) = SO_0(\eta) \times \mathbb{R}^+$, $v = (v_a) \in \mathbb{R}^{n*}$. It follows that H is isomorphic to the semi-direct product $CO_0(\eta) \times \mathbb{R}^{n*}$ with the multiplication law

$$(h^a_b, v_a) (k^a_b, w_a) = (h^a_r k^r_b, v_r k^r_a + w_a),$$

where $h^a_b = \exp(\sigma) \Lambda^a_b$, with $\exp(\sigma) \in \mathbb{R}^+$ and $\Lambda \in CO_0(\eta)$. With (h^a_b, v_a) written as $(\theta, \Lambda^a_b, v_a)$, one can easily verify that the following formula defines a representation R of H on $\mathbb{R}^{n+2} = \mathbb{R}^n \oplus \mathbb{R}^2$:

$$R(\theta, \Lambda, v) = \begin{pmatrix} \Lambda^r_s & \eta^{rd} v_s & \eta^{rs} v_s \\ -v_r & \frac{1+\theta^2-v^2}{2\theta} & -\frac{1-\theta^2+v^2}{2\theta} \\ \frac{v_r}{\theta} & -\frac{1-\theta^2-v^2}{1\theta} & \frac{1+\theta^2+v^2}{2\theta} \end{pmatrix}.$$

With $S = \begin{pmatrix} \eta & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$ we then have $R(\theta, \Lambda, v)^t S R(\theta, \Lambda, v) = S$, therefore the representation R realizes H as a subgroup of the group $G = SO_0(p+1, q+1)$.

The part of G that is missing in H is the translation group given by the following $SO_0(p+1, q+1)$ matrices $T(a)$, $a \in \mathbb{R}^n$:

$$T(a) = \begin{pmatrix} \delta_s^r & -a^r & a^r \\ \eta_{rs}a^s & 1 - a^2/2 & a^2/2 \\ \eta_{rs}a^s & -a^2/2 & 1 + a^2/2 \end{pmatrix}, \quad (32)$$

- Cf. section 5.2. The Lie algebra generators $so(p+1, q+a)$ take now the following form:

$$D = \frac{dD(\exp(\sigma), E, 0)}{d\sigma} \Big|_{\sigma=0} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$\frac{1}{2}\omega^r_s M^s_r = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \omega^r_s & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$v_r K^r = \begin{pmatrix} 0 & \eta^{rs}v_s & \eta^{rs}v_s \\ -v_r & 0 & 0 \\ v_r & 0 & 0 \end{pmatrix}$$

$$w^r P_r = \begin{pmatrix} 0 & -w^r & w^r \\ \eta_{rs}w^s & 0 & 0 \\ \eta_{rs}w^s & 0 & 0 \end{pmatrix}$$

8.4 The enlarged conformal bundle and the normal Cartan connection

With H being a subgroup of G , as above, we can build now the associated bundle $\tilde{P}_C^2(M) = P_C^2(M) \times_H G$, which is a principal G -bundle (cf. e.g., [39, p. 4] and references therein). If $n = p+q \geq 3$, then this new bundle is naturally equipped with a principal connection, the normal Cartan connection, which can be described as follows.

Let g be a metric in the conformal class C , let e_a be an (local) orthonormal frame of g , and R its curvature tensor. Then, in a coordinate system x^μ , the covariant derivative $\nabla_\mu Z$ of a section Z of the associated vector bundle $\tilde{P} \times_R E^{p+1, q+1}$ is given by the following expression - cf. e.g., [37, Ch. 4.4], [39, p. 14], [23, p. 196] :

$$\nabla_\mu Z = \partial_\mu Z + \Gamma_\mu Z,$$

with

$$\Gamma_{\mu} = \frac{1}{2}\Gamma_{\mu}^{rs}M_r^s + \frac{1}{n-2}\left(R_{\mu\sigma} - \frac{1}{2(n-1)}Rg_{\mu\sigma}\right)K^{\sigma} - P_{\mu},$$

where $K^{\mu} = e_r^{\mu}$, and $P_{\mu} = e_{\mu}^r P_r$.

In a natural way we can then build the associate bundle $\tilde{P} \times_G E^{p+1,q+1}$ with $E^{p+1,q+1}$ as a typical fibre, and we can construct the projective quadric \tilde{M}_x at each point $x \in M$.

Now, suppose M is connected and simply connected and the conformal structure is flat. In this case we can choose (cf. [37, Ch. I.2]) $g_{\mu\nu} = \eta_{\mu\nu}$. The covariant derivative $\nabla_{\mu}Z$ reduces in this case to

$$\nabla_{\mu}Z = \partial_{\mu}Z - P_{\mu}Z.$$

In an adapted coordinate system x^{μ} we choose the ‘‘origin’’ of the ‘‘compactified tangent space’’ to correspond to the point $(0, \frac{1}{2}, -\frac{1}{2})$ of $E^{p+1,q+1}$. Connecting the point $x \in M$ with $0 \in M$ by the path $x(t) = (1-t)x$ we can then transport parallelly the origin $(0, \frac{1}{2}, -\frac{1}{2})$ at to the point $0 \in M$. The parallel transport rule gives us

$$0 = DZ(x(t))/dt = dZ(x(t))/dt - dx^{\mu}/dt P_{\mu}Z(x(x(t))),$$

or, in our case, $dZ/dt = -x^{\mu}P_{\mu}Z$, which solves to

$$Z(1) = \exp(x^{\mu}P_{\mu})Z(0),$$

or, applying Eq. (32):

$$Z(1) = (x, (1-x^2)/2), -(1+x^2)/2),$$

which is nothing but the standard embedding (9).

9 Concluding remarks

This paper has provided a mathematical analysis of algebraic and geometrical aspects of the Minkowski space compactification. Some omissions, faulty reasoning and lack of precision in the existing literature dealing with this subject has been pointed out and analyzed in some detail. In addition to the standard compactification by adding a ‘‘light cone and a 2-sphere at infinity’’ also its double covering isomorphic to $U(1) \times SU(2)$ has been discussed. A

pictorial representation has been proposed and the corresponding “Penrose diagrams” have been derived. The role of the conformal inversion and the representation of null geodesics has been touched upon as well. Applications to flat conformal structures, including the normal Cartan connection and conformal development has been discussed in some detail. In appendix A a detailed discussion of the spaces of null lines in a general case of a pseudo-Hermitian space $H_{p,q}$ has been given.

10 Acknowledgments

Thanks are due to Pierre Anglès and Rafał Abłamowicz for their critical reading parts of the manuscript and for the discussion, and to Don Marolf for pointing out a possible usefulness of adding the analysis of the $IO(3,1)$ action, also to Alexander Levichev for useful suggestions. Special thanks are also due to Paweł Nurowski for sharing with me some of his knowledge and for pointing to me the question discussed in appendix A. I am indebted to Nikolay M. Nikolov for sending me some of his papers and for discussion.

Appendices

A Killing vector fields for the left action of $U(2)$ on itself

A.1 The problem

We take the group $U(2)$ in the standard matrix form. It has the manifold structure of $(S^1 \times S^3)/Z_2$ - the same as the compactified Minkowski space.

Now, let ω be the Maurer-Cartan form of $U(2)$, a 2×2 matrix of one-forms. Taking the determinant of ω with understanding that the multiplication of one-forms is to be understood as a symmetrized tensor product, we obtain a symmetric bilinear form $g = \det(\omega)$. This form is non-degenerate of Lorentzian signature and is conformal to the flat Minkowski metric under the standard identification of $U(2)$ as the compactification of the Minkowski space M . The metric g obtained this way is, by its very construction, invariant under the left action of $U(2)$ on itself. Therefore the left action of $U(2)$

on itself leads to conformal transformations of M .

Precisely which subgroup of the conformal group corresponds to this left action of $U(2)$ on itself?

A.2 The solution

It is well known that the group $SU(2, 2)$ acts by conformal automorphisms on the compactified Minkowski space (see e.g., [9]). The group $U(2, 2)$ consists of block matrices $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ with entries A, B, C, D which are 2×2 complex matrices satisfying the relations $A^*A - C^*C = D^*D - B^*B = E$ and $A^*B - C^*D = 0$. Its action on $U(2)$ is given by the fractional linear transformations:

$$U \mapsto U' = (AU + B)(CU + D)^{-1}, \quad (33)$$

with $CZ + D$ being automatically invertible for $U \in U(2)$. By specifying $B = C = 0$, $D = E$, we see that A is in $U(2)$. Therefore the left action of $U(2)$ on itself is a particular case of the linear fractional transformations as above.

In order to describe these transformations in the Minkowski space, we can use the Cayley transform as in [1]. Or, we can inverse Cayley-transform the matrices of $U(2, 2)$ and act on the Minkowski space represented by hermitian 2×2 matrices in the standard form: $X = x^\mu \sigma_\mu$, where

$$\sigma_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

The action of $U(2, 2)$ is still described by fractional linear transformations

$$X \mapsto X' = (RX + S)(TX + Q)^{-1}, \quad (34)$$

where (cf. [22, (2.16)]) $A = \frac{1}{2}(R + iS - iT + Q)$, $B = \frac{1}{2}(-R + iS + iT + Q)$, $C = \frac{1}{2}(-R - iS - iT + Q)$, $D = \frac{1}{2}(R - iS + iT + Q)$. With $B = C = 0$ and $D = E$ we easily find that $R = \frac{1}{2}(A + E)$, $Q = \frac{1}{2}(A + E)$, $S = -i\frac{1}{2}(A - E)$, $T = i\frac{1}{2}(A - E)$. Consider now a one-parameter subgroup $A(\tau)$ of $U(2)$. By differentiating the equation

$$X(\tau) = (R(\tau)X + S(\tau))(T(\tau)X + Q(\tau))^{-1}$$

at $\tau = 0$, and putting $A(0) = E$, $\dot{A}(0) = i\sigma$ we obtain:

$$\dot{X} = \frac{i}{2}(\sigma X - X\sigma) + \frac{1}{2}\sigma + \frac{1}{2}X\sigma X.$$

Denoting by Z_μ the vector fields corresponding to $\sigma = \sigma_\mu$ we easily find their components using the simple algebra:

$$(Z_\mu)^\nu = \frac{1}{2} \text{tr}(\dot{X} \sigma_\nu).$$

The result is as follows:

$$Z_0 = \frac{1}{2} (1 + t^2 + x^2 + y^2 + z^2) \partial_t + t(x\partial_x + y\partial_y + z\partial_z),$$

$$Z_1 = tx \partial_t + \frac{1}{2} (1 + t^2 + x^2 - y^2 - z^2) \partial_x + (xy + z)\partial_y + (xz - y)\partial_z$$

$$Z_2 = ty \partial_t + (xy - z)\partial_x + \frac{1}{2} (1 + t^2 - x^2 + y^2 - z^2) \partial_y + (xz + y)\partial_z$$

$$Z_3 = tz \partial_t + (xz + y)\partial_x + (yz - x)\partial_y + \frac{1}{2} (1 + t^2 - x^2 - y^2 + z^2) \partial_z.$$

We can now compare these vector fields with the formulas for the standard generators P_μ , K_μ , $M_{\mu\nu}$ of the conformal group as given, for instance, in [40]:

$$\begin{aligned} P_\mu &= -\partial_\mu \\ M_{\mu\nu} &= x_\mu \partial_\nu - x_\nu \partial_\mu \\ K_\mu &= -2x_\mu (x^\nu \partial_\nu) + x^2 \partial_\mu \end{aligned}$$

By an easy calculation we find:

$$Z_0 = \frac{1}{2}(K_0 + P_0), \quad Z_i = \frac{1}{2}(K_i - P_i) + L_i, \quad (i = 1, 2, 3),$$

where $L_i = \epsilon_{ijk} M_{jk}$.

B Comments on projective quadrics subordinate to pseudo-Hermitian spaces

In this section we present a self-contained generalization of some of our constructions to a signature (p, q) . The presentation below is the result of discussing with the Author the reasoning in [23, pp. 209-212]. Pierre Anglès subsequently published a corrected derivation [41], which gives, by a different

method, the results presented below. This appendix contains the material referred to as *Comments on projective quadrics subordinate to pseud-Hermitian spaces* in the References section of [41].

Let \mathbb{C} be the field of complex numbers, and let C^* be the multiplicative group of complex numbers different from zero. Using the polar decomposition we can write $C^* = \mathbb{R}^+ \times U(1)$, where \mathbb{R}^+ is the multiplicative group of positive real numbers and $U(1)$ is the circle group.

Let V be a complex vector space of finite dimension n , equipped with a regular pseudo-hermitian form (x, y) of signature (p, q) , $p + q = n$, $p, q \geq 1$. and let Q be the isotropic cone minus the origin:

$$Q = \{x \in V : (x, x) = 0, x \neq 0\}.$$

Q is a real manifold of (real) dimension $2n - 1$, and we denote by \tilde{Q} the quotient manifold $\tilde{Q} = Q/\mathbb{C}^*$. Its elements are the equivalence classes: $\tilde{Q} = \{\{cx : c \in \mathbb{C}^*\}, x \in Q\}$. \tilde{Q} is a real submanifold of the complex projective space $P(V)$. We denote by P the canonical projection $P : V \rightarrow P(V)$.

The projection P can be implemented in two steps: first taking the quotient with respect to \mathbb{R}^+ to obtain $Q' = Q/\mathbb{R}^+$, then quotienting Q' by $U(1)$ to obtain \tilde{Q} . We denote the corresponding projections P' and π respectively. Thus we have $P = \pi \circ P'$, and $\tilde{Q} = Q'/U(1)$. Q' and \tilde{Q} are real compact manifolds of dimensions $2n - 2$ and $2n - 3$ respectively.²⁰

Let $T_x Q$ be the tangent space at $x \in Q$. Then $T_x Q$ can be identified with a real vector subspace of V as follows. If $\mathbb{R} \ni t \mapsto x(t) \in Q$ is a path with $x(0) = x$, then $(x(t), x(t)) = 0 \forall t$. Denoting by $X = \left. \frac{dx(t)}{dt} \right|_{t=0}$ the tangent vector at x , and using the Leibniz rule, we get $(X, x) + (x, X) = 0$, or

$$X \in T_x Q \text{ if and only if } \operatorname{Re}((X, x)) = 0.$$

Let $T_x^{\mathbb{C}} Q = x^{\perp}$ be the subspace of $T_x Q$ defined by the condition $(X, x) = 0$. Then $T_x^{\mathbb{C}} Q$ is a hyperplane in $T_x Q$. In fact, while $T_x Q$ is only a real vector space, $T_x^{\mathbb{C}} Q$ carries the structure of a complex space.

Let η_{jk} be the diagonal matrix $\eta_{jk} = \delta_{jk}$ for $j, k = 1, \dots, p$, $\eta_{jk} = -\delta_{jk}$ for $j, k = p + 1, \dots, p + q$. Let $H_{p,q}$ be the standard pseudo-hermitian space \mathbb{C}^{p+q} equipped with the scalar product

$$f(u, v) = \sum_{j,k=1}^n \eta_{jk} u^j \bar{v}^k.$$

²⁰In fact, we will see that Q' is diffeomorphic to the product of two odd-dimensional spheres $S^{2p-1} \times S^{2q-1}$.

$H_{p,q}$ is the direct sum of two subspaces $H_{p,q} = H_{p,q}^+ \oplus H_{p,q}^-$ spanned by the first p (resp. last q) vectors of the standard basis. Every orthonormal basis $\{e_j\}$, $(e_j, e_k) = \eta_{jk}$ in V determines an isometry $\phi_e : H_{p,q} \rightarrow V$ and determines an orthogonal direct sum decomposition $V = \phi_e(H_{p,q}^+) \oplus \phi_e(H_{p,q}^-)$ into a positive and a negative subspace. We call such a decomposition a “split”. We denote by \mathcal{S} the set of all splits of V . Then \mathcal{S} is a homogeneous space (in fact, it is a Kähler manifold) for the unitary group $U(V)$, isomorphic to $U(p, q)/(U(p) \times U(q))$.

B.1 The topology of $Q' = P'(Q)$.

Let $\{e_j\}$ be an orthonormal basis in V , so that we can identify V with $H^{p,q}$. The equation of Q becomes then

$$\sum_{j=1}^p |z^j|^2 = \sum_{j=p+1}^{p+q} |z^j|^2 \neq 0.$$

Consider the submanifold S_1 of $H^{p,q}$ defined by the formula

$$\sum_{j=1}^p |z^j|^2 = \sum_{j=p+1}^{p+q} |z^j|^2 = 1. \quad (35)$$

Then S_1 is the product of two spheres $S_1 = S^{2p-1} \times S^{2q-1}$ and the projection P' restricted to S_1 is a diffeomorphism from S_1 to $Q' = P'(Q)$. The representation of Q' as a product of two spheres will, in general, depend on the choice of the orthonormal basis, more specifically: on the split determined by the basis.

Somewhat more generally, let $s \in \mathcal{S}$ be a split, so that V is decomposed into a direct (orthogonal) sum $V = V_+ \oplus V_-$ of positive and negative subspaces. Defining $\|x\| = (x, x)$ on V_+ , and $\|x\| = -(x, x)$ on V_- , each isotropic vector $x \in Q$ decomposes into a sum $x = x_+ + x_-$, with $x_+ \in V_+$, $x_- \in V_-$, and $\|x_+\| = \|x_-\| = R(x) > 0$. Rescaling $x \mapsto x/R(x)$ we get the unique representative of the equivalence class \mathbb{R}^+x of x with $R(x) = 1$. In other words, if we define

$$Q_s = \{x \in Q : \|x_+\| = \|x_-\| = 1\},$$

then Q_s defines a global cross section of the projection $P' : Q \rightarrow Q'$, and a diffeomorphism of Q' onto the product of the two unit spheres, one in V_+ and one in V_- .

B.2 The conformal structure of Q'

Let a be a point of Q' and let x be an isotropic vector in Q with $P(x) = a$. The tangent space $T_x Q$ is equipped with the (real) bilinear form $f_x(X, Y) = \operatorname{Re}((X, Y))$. Notice that the vector x itself can be considered as an element of $T_x Q$, and that the line $\mathbb{R}x$ is the radical of the bilinear form f_x , and is the kernel of the tangent map $(dP')_x$:

$$\mathbb{R} = \{y \in T_x Q : f_x(y, z) = 0 \forall z \in T_x Q\} = \{y \in T_x Q : (dP')_x(y) = 0\}.$$

It follows that the form f_x induces a regular bilinear form, which we denote g_x on the tangent space $T_a Q'$.

Lemma 5. *With the notation as above, if $\lambda > 0$ then $g_{\lambda x} = \lambda^2 g_x$.*

Proof. Let $a_1(t), a_2(t)$, $a_1(0) = a_2(0) = a$, be two paths in Q' with tangent vectors $\dot{a}_1(0)$ and $\dot{a}_2(0)$ respectively. Let x_1, x_2 be the lifts: $P'(x_1(t)) = a_1(t)$, $P'(x_2(t)) = a_2(t)$, $x_1(0) = x_2(0) = x$. Then, according to the definition of g_x , we have that²¹

$$g_x(\dot{a}_1(0), \dot{a}_2(0)) = f_x(\dot{x}_1(0), \dot{x}_2(0)).$$

Let $\lambda > 0$, then $x'_1(t) = \lambda x_1(t)$ and $x'_2(t) = \lambda x_2(t)$ are lifts through λx of $a_1(t)$ and $a_2(t)$ respectively, with tangent vectors $\lambda \dot{x}_1$, $\lambda \dot{x}_2$. Thus $g_{\lambda x}(\dot{a}_1(0), \dot{a}_2(0)) = \lambda^2 g_x(\dot{a}_1(0), \dot{a}_2(0))$. \square

It follows from the above lemma that what is independent of the choice of x in $P'^{-1}(a)$ is the conformal class of g_x and, in particular, the signature, which, by construction, is $(2p - 1, 2q - 1)$.²²

In order to proceed further on notice that we have the following, easy to prove, lemma:

Lemma 6. *Given a vector $x \in Q$ there exists an orthonormal basis e such that $x = e_1 + e_n$.*

²¹Notice that this expression does not depend on the choice of the lifts, the reason being that vectors tangent to two lifts will differ by vectors in the kernel of $(dP')_x$, that is in $\mathbb{R}x$, which is orthogonal to all vectors in $T_x Q$.

²²The fact that the conformal class of the induced metric on Q_s does not depend on the choice of the split s is by no means evident if the two induced metrics are computed using two different orthonormal bases related by a general $U(p, q)$ transformation and then compared.

Proof. Take any orthonormal basis $\{e'_i\}$. The vector $x' = e'_1 + e'_n$ is isotropic. We know that the automorphism group of V acts transitively on isotropic lines (see e.g., [42, p. 74, Corollaire 2]). Let U be any automorphism of V with the property $x = Ux'$, and let $e_i = Ue'_i$. Then $\{e_i\}$ is an orthonormal basis of V and $x = e_1 + e_n$. \square

Each tangent space $T_x(Q)$ is also equipped with a skew-symmetric bilinear form F_x defined by

$$F_x(X, Y) = \text{Im}(X, Y), \quad X, Y \in T_x(Q).$$

However the form F_x does not descend to the quotient $Q' = Q/\mathbb{R}x$ because, owing to the fact that, for instance, ie_1 is in $T_x(Q)$ but $F_x(x, ie_1) = 1$, we find that x is not in the radical of F_x . But F_x , **when restricted to x^\perp** , does descend to a skew-symmetric bilinear form on $(dP')_x(x^\perp)$.

Because of the lemma above it is instructive to consider first the case of $p = 1, q = 1$.

B.3 The case of the signature (1, 1).

With $x \in Q$, let (e_1, e_2) be the orthonormal basis $(e_1, e_1) = 1, (e_2, e_2) = -1$, with $x = e_1 + e_2$, and let $\{f_1, f_2, f_3, f_4\}$ $f_1 = e_1 + e_2 = x, f_2 = i(e_1 + e_2), f_3 = i(e_1 - e_2), f_4 = (e_1 - e_2)$. The tangent space $T_x Q$ is spanned by the vectors $\{f_1, f_2, f_3\}$, the complex orthogonal space x^\perp is spanned by the vectors $\{f_1, f_2\}$. Let $a = P'(x)$. Notice that $(dP')_x(f_1) = 0$, while $(dP')_x$ is a bijection from the plane spanned by $\{f_2, f_3\}$ onto $T_a Q'$. Denoting $\epsilon_1 = (dP')_x(ie_1), \epsilon_2 = (dP')_x(ie_2)$, the vectors ϵ_1, ϵ_2 form an orthonormal basis in $T_a Q'$ for the induced bilinear form g_x :

$$g_x(\epsilon_1, \epsilon_1) = -g_x(\epsilon_2, \epsilon_2) = 1, \quad g_x(\epsilon_1, \epsilon_2) = 0.$$

Q' is now the torus $S^1 \times S^1$ given by the formula (35), now becoming:

$$|z^1|^2 = |z^2|^2 = 1.$$

Writing $z^1 = \cos(\phi_1) + i \sin(\phi_1), z^2 = \cos(\phi_2) + i \sin(\phi_2)$, the pseudo Riemannian metric g_x of Q' , when expressed in the natural torus coordinates ϕ_1, ϕ_2 is diagonal $g_x = \text{diag}(1, -1)$. The action $z \mapsto \exp(i\phi)z$ of $U(1)$ on Q translates to the action $(\phi_1, \phi_2) \mapsto (\phi_1 + \phi, \phi_2 + \phi)$ on the torus. The tangent vector to the orbit of this action at x is f_2 that projects onto $\epsilon_1 + \epsilon_2$ at $T_a Q'$. Taking

the quotient of Q' by this action we get \tilde{Q} as the circle $\tilde{Q} = Q/\mathbb{C}x = S^1$. The image of $x^\perp = \mathbb{C}x$ by $(dP)_x$ consists of one point - the zero vector. Since $\epsilon_1 + \epsilon_2$ is a null vector for the metric g_x , there is no distinguished subspace transversal to the fiber, therefore no metric whatsoever is generated by dP on \tilde{Q} .

B.4 The structure of \tilde{Q}

Given a split $s \in \mathcal{S}$, let $V = V_+ \oplus V_-$ be the corresponding decomposition V . Every vector $x \in Q$ can be then uniquely represented as $x = x_+ + x_-$, so that $(x_+, x_+) = (x_-, x_-)$. With

$$Q_s = \{x \in Q : (x_+, x_+) = (x_-, x_-) = 1\},$$

the map $P : Q \rightarrow Q'$, when restricted to Q_s , becomes a diffeomorphism. The $U(1)$ action $x \mapsto cx$, $|c| = 1$ leaves Q_s invariant. (Q', π) is a $U(1)$ principal fibre bundle over \tilde{Q} . Given a split s , Q' is endowed with the pseudo-Riemannian metric g_s that is automatically $U(1)$ -invariant.

Given a non-degenerate pseudo-Riemannian metric on a principal bundle, the standard method of obtaining the metric on the base space is by taking the orthogonal complement to the fibers. This method works when the orthogonal complement is transversal to the fibers. Yet in our case the vectors tangent to the fibers are isotropic, therefore the orthogonal complement is not transversal to the fibers. Nevertheless, we can obtain a natural, though degenerate, scalar product g on the cotangent bundle of \tilde{Q} . as follows:

Let \mathcal{S} be as split, let $a \in \tilde{Q}$, and let ω, ω' be two one-forms in the cotangent space $T_a^*\tilde{Q}$. Let $b \in Q'$ be a point in the fibre $\pi^{-1}(a)$. The pullbacks $\pi_*\omega, \pi_*\omega'$ are invariant one-forms defined at the points of the fibre $\pi^{-1}(a)$. We can therefore calculate the scalar product $g_x^*(\pi_*\omega, \pi_*\omega')$, at any point of the fibre, and, owing to the fact that the forms and the metric are invariant, the result is independent of the chosen point. Since the scalar products corresponding to different choices of x differ only by a scale factor, the same is true about the induced contravariant symmetric scalar product on the cotangent bundle of \tilde{Q} . The scalar product so obtained is degenerate. Indeed, any form that vanishes on the image $(dP')_x(x^\perp)$ is in the radical of g_x^* .

There is another way of looking at this construction. Let W_1 be a subspace of a real vector space W , and let f_1 be a non-degenerate

symmetric bilinear form on W_1 . Let $\iota : W_1 \rightarrow W$ be the canonical inclusion map, and let $\iota^* : W^* \rightarrow W_1^*$ be its dual. The bilinear form f_1 can be considered as a map $f_1 : W_1 \rightarrow W_1^*$, and, since we assume it to be non-degenerate, there exists the inverse $f_1^* : W_1^* \rightarrow W_1$. We can then define $f^* : W^* \rightarrow W$ by

$$f^* = \iota \circ f_1^* \circ \iota^*.$$

The map f^* can now be considered as a bilinear form on W^* and it is easy to see that, by construction, it is symmetric. Moreover, its radical consists of the forms $\omega \in W^*$ that vanish on the image $\iota(W_1)$.

B.5 \tilde{Q} as a compactification of $\mathbb{R} \times H_{p-1,q-1}$

Let $a, b \in \tilde{Q}$.

Definition 1. We write $a \perp b$ if and only if $a = P(x)$, $b = P(y)$, where $x, y \in Q$, and $(x, y) = 0$. Given $a \in \tilde{Q}$ we define

$$a^\perp = \{b \in \tilde{Q} : a \perp b\}.$$

It can be seen that a^\perp is a closed subset of \tilde{Q} .

Let us fix $a \in \tilde{Q}$, and let $x \in Q$ be such that $P(x) = a$. We recall that x^\perp is a complex vector subspace of V that carries a degenerate sesquilinear form inherited from the scalar product of V , with radical $\mathbb{C}x$. Therefore the quotient space

$$M \stackrel{\text{df}}{=} x^\perp / \mathbb{C}x$$

carries the pseudo-Hermitian form of signature $(p-1, q-1)$. We can realize M as follows: choose $u \in Q$ such that $(u, x) = 1$.²³ Let M_u be the orthogonal complement of $\{x, u\}$ in V . Then the scalar product of V restricted to M_u is of signature $(p-1, q-1)$, we evidently have $M_u \subset x^\perp$, and the projection $x^\perp \rightarrow x^\perp / \mathbb{C}x$ restricted to M_u is easily seen to be a bijection.

We will construct now a bijection κ from $\mathbb{R} \times M_u$ onto $\tilde{Q} \setminus a^\perp \subset \tilde{Q}$. Given $r \in \mathbb{R}$, $y \in M_u$ define

$$\kappa_0(r, y) = y + u + \left(-\frac{1}{2}(y, y) + ri \right) x.$$

²³Such a choice is always possible, for instance, by using Lemma 2, we can set $x = e_1 + e_n$, $u = (e_1 - e_n)/2$.

Notice that the coefficient in front of x has the imaginary part r . It is easy to see that, automatically, $\kappa_0(r, y) \in Q$ and also $(x, \kappa_0(r, y)) = 1$.

We define now $\kappa = P \circ \kappa_0$. It is easy to check that κ is injective. It remains to show that it is a surjection from $\mathbb{R} \times M_u$ onto $\tilde{Q} \setminus a^\perp$. Given $b \in \tilde{Q} \setminus a^\perp$, let z' be any point in $P^{-1}(b)$. Then, since b is not in a^\perp , we have that $(z', x) = a \neq 0$. Taking $z = z'/a$, we still have $P(z) = b$, but now $(z, x) = 1$. Now, z can be uniquely written in the form $z = y + \alpha u + \beta x$, where $y \in M_u$, $\alpha, \beta \in \mathbb{C}$. From $(z, x) = 1$ we find that $\alpha = 1$, and from $(z, z) = 0$ we get that $\text{Re}(\beta) = -\frac{1}{2}(y, y)$. Putting $r = \text{Im}(\beta)$ we get $b = \kappa(r, y)$.

B.5.1 The structure of a^\perp .

Proposition 3. *With the notation as above,*

$$a^\perp \cong \{1\} \cup \mathbb{R} \times (S^{p-2} \times S^{q-2})/S^1.$$

Proof. Let $\{e_j\}$ be an orthonormal basis such that $x = e_1 + e_n$. Then any vector $y \in x^\perp$ is of the form

$$y = \alpha x + \sum_{j=2}^{n-1} \alpha^j e_j.$$

Such a vector y is in Q if and only if $\sum_{j=2}^p |\alpha^j|^2 = \sum_{j=p+1}^{n-1} |\alpha_j|^2$. If all α^j are zero, then, necessarily, $\alpha \neq 0$, and we can choose a unique representative of the equivalence class with $\alpha = 1$. This gives us the point $\{1\}$. If at least one of the α^j is non-zero, then $\sum_{j=2}^p |\alpha^j|^2 = \sum_{j=p+1}^{n-1} |\alpha_j|^2 \neq 0$ and we can use the freedom of real scaling to get $\sum_{j=2}^p |\alpha^j|^2 = \sum_{j=p+1}^{n-1} |\alpha_j|^2 = 1$. The remaining freedom of $U(1)$ gives us $(S^{p-2} \times S^{q-2})/S^1$. The α coefficient remains still arbitrary, thus the result follows. \square

References

- [1] Armin Uhlmann, *The Closure of Minkowski Space*, Acta Physica Polonica, Vol. XXIV, Fasc. 2(8), pp. 295–296 (1963)
- [2] R. Penrose and W. Rindler, *Spinors and Space-Time, Vol. 2 – Spinor and Twistor Methods in Space-Time Geometry*, Cambridge University Press, Cambridge, England, 1984

- [3] S. A. Huggett and K. P. Tod, *An Introduction to Twistor Theory*, Cambridge University Press (1994)
- [4] John K. Beem, Paul E. Ehrlich, Kevin L. Easley, *Global Lorentzian Geometry, Second Edition*, Marcel Dekker, Inc., New York (1996)
- [5] José L. Flores, *The Causal Boundary of Spacetimes Revisited*, Commun. Math. Phys. **276**, pp. 611–643 (2007)
- [6] David E. Lerner, *Global Properties of Massless Free Fields* Commun. Math. Phys. **55**, pp. 179–182 (1977)
- [7] Claude Chevalley, *Theory of Lie Groups I*, Princeton Mathematical Series, no. 8, Princeton University Press, (1946), Chapter I, §X, Proposition 6
- [8] A. V. Levichev, *Parallelizations of Chronometric Bundles Based on the Subgroup $U(2)$* , Izvestia RAEN, ser.MMMIU, 10 (2006), n.1-2, pp. 51–61, (in Russian)
- [9] W. Kopczyński and L. S. Woronowicz, *A geometrical approach to the twistor formalism*, Rep. Math. Phys. Vol **2**, pp. 35–51 (1971)
- [10] Leszek Z. Stolarczyk, *The Hodge Operator in Fermionic Fock Space*, Collect. Czech. Chem. Commun. **70**(2005), pp. 979–1016
- [11] René Deheuvels, *Formes quadratiques et groupes classiques*, Presse Universitaires de France, Paris (1981)
- [12] Aubert Daigneault, *Irving Segal's axiomatization of spacetime and its cosmological consequences*, Preprint <http://arxiv.org/abs/gr-qc/0512059>
- [13] B. G. Schmidt, *A New Definition of Conformal and Projective Infinity of Space–Times*, Commun. math. Phys. **36** (1974), pp. 73–90
- [14] Maks A. Akivis, Vladislav V. Goldberg, *Conformal Differential Geometry and its Generalizations*, A Wiley Interscience Publications, New York (1996)

- [15] N. M. Todorov, I. T. Todorov, *Conformal Quantum Field Theory in Two and Four Dimensions*, Vienna, Preprint ESI 1155 (2002), <http://www.esi.ac.at/preprints/esi1155.ps>
- [16] Nikolay M. Nikolov, *Rationality of Conformally Invariant Local Correlation Functions on Compactified Minkowski Space*, Commun. Math. Phys. **218** (2001), pp. 417–436
- [17] Nikolay M. Nikolov, *Vertex Algebras in Higher Dimensions and Globally Conformal Invariant Quantum Field Theory*, Commun. Math. Phys. **253** (2005), pp. 283–322
- [18] Roger Penrose, *Conformal treatment of Infinity*, in “Relativity, groups and topology”, Lectures delivered at Les Houches during the 1963 session of the Summer School of Theoretical Physics, University of Grenoble, ed. DeWitt, Gordon and Brach, New York (1964), pp. 563–584
- [19] C. W. Misner, K. S. Thorne, J. A. Wheeler, *Gravitation*, Freeman and Co., New York (1973)
- [20] S. W. Hawking, G. F. R. Ellis, *The large scale structure of space-time*, Cambridge University Press, Cambridge (1976)
- [21] Roger Penrose, *The Light Cone at Infinity*, in “Relativistic Theories of Gravitation”, ed. L. Infeld, Pergamon Press, Oxford (1964), pp. 369–373
- [22] W. Rühl, *Distributions on Minkowski Space and their Connection with Analytic Representations of the Conformal Group*, Commun. math. Phys., **27** (1972), pp. 53-86
- [23] Pierre Angles, *Conformal Groups in Geometry and Spin Structures*, Birkhauser, Progress in Mathematical Physics, Vol. 50 (2008)
- [24] Roger Penrose, *Structure of space-time*, in Battelle Rencontres, ed. by C. M. DeWitt, J. A. Wheeler, Benjamin, New York (1969), pp. 121–235,
- [25] Roger Penrose, *Twistor Algebra*, J. Math. Phys., **8** (1967), pp. 345–366
- [26] N. H. Kuiper, *On conformally flat spaces in the large*, Ann. of Math. **50** (1949), pp. 916–924

- [27] R. Penrose, *Zero Rest-Mass Fields Including Gravitation: Asymptotic Behaviour* Proc. R. Soc. London **284** (1965), pp. 159–203
- [28] Anil Zenginoğlu, *Hyperboloidal foliations and scri-fixing*, <http://arxiv.org/abs/0712.4333>
- [29] Robert Geroch, *Asymptotic Structure of Space-Time*, in “Asymptotic Structure of Space-Time”, ed. F. Esposito and L. Witten, Plenum Press (1977), pp. 1–105
- [30] M. Chisholm, *Such Silver Currents. The Story of William and Lucy Clifford 1845–1929*, The Lutterworth Press, Cambridge (2002)
- [31] Roger Penrose, *The Road to Reality*, Jonathan Cape, London (2004)
- [32] Werner Greub, *Multilinear Algebra*, Springer (1978)
- [33] John. C. Baez, Irving E. Segal, Zhengfang Zhou, *Introduction to Algebraic and Constructive Quantum Field Theory*, Princeton University Press (1992)
- [34] W. D. Curtis, F. R. Miller, *Differential Manifolds and Theoretical Physics*, Academic Press, New York (1985)
- [35] Fecko, M.: *Differential Geometry and Lie Groups for Physicists*, (Cambridge University Press, Cambridge (2006)
- [36] David E. Lerner, *Twistors and induced representations of $SU(2,2)$* , J. Math. Phys. **18** (1977), pp. 1812–1817
- [37] Shoshichi Kobayashi, *Transformation Groups in Differential Geometry*, Springer (1972)
- [38] A. Jadczyk, *Born’s Reciprocity in the Conformal Domain*, in Z. Oziewicz et al (eds.), *Spinors, Twistors, Clifford Algebras and Quantum Deformations*, Kluwer Academic Publishers (1993), pp. 129–140
- [39] Felipe Leitner, *Twistor Spinors and Normal Cartan Connections in Conformal Geometries*, <ftp://ftp-sfb288.math.tu-berlin.de/pub/Preprints/preprint471.ps.gz>
- [40] J. Mickelsson, J. Niederle, *Conformally invariant field equations*, Ann. de l’I.H.P., section A, tome 23, no 3 (1975), p. 277–295

- [41] Pierre Anglès, *Study of some fundamental projective quadrics associated with a standard pseudo-Hermitian space $H_{p,q}$* , To appear in *Advances in Applied Clifford Algebras*
- [42] Bourbaki Nicolas, *Éléments de Mathématique. Algèbre, Chapitre 9*, Springer (1981)




Figure 3: Pictorial representation of the conformal infinity with one dimension skipped. Double light cone at infinity with endpoints identified. While topologically correct this representation is misleading as it suggests non differentiability at the base, where the two half-cones meet.




Figure 4: Conformal infinity of Minkowski space from [4, p. 178]. The meaning of this picture is quite different from the one in Fig. 3, where the points i^0, i^+, i^- are identified. The 2-sphere indicated in the middle of this picture is just one point i^0 and **not** the true 2-sphere of Fig. 3




Figure 5: Chronological boundary for \mathbb{L}^{n+1} - Figure. 2 from [5]




Figure 6: Differentiably correct pictorial representation of the conformal infinity with one dimension skipped. A torus squeezed to a point $I^+ = I^- = I^0$ at $\psi = 0$. All null geodesics described in this section pass through this point.




Figure 7: Pictorial representation of the double covering conformal infinity. Double double light cone. Points connected by a dashed line are, in fact, a one point. This representation is also topologically correct but differentiably misleading.




Figure 8: Pictorial representation of the double covering conformal infinity.
A pair of tori squeezed at a common point.